

HST STIS SPECTROSCOPY OF THE TRIPLE NUCLEUS OF M31: TWO NESTED DISKS IN KEPLERIAN ROTATION AROUND A SUPERMASSIVE BLACK HOLE

RALF BENDER,^{1,2,3} JOHN KORMENDY,⁴ GARY BOWER,⁵ RICHARD GREEN,⁶ JENS THOMAS,^{1,2} ANTHONY C. DANKS,⁷
THEODORE GULL,⁸ J. B. HUTCHINGS,⁹ C. L. JOSEPH,¹⁰ M. E. KAISER,¹¹ TOD R. LAUER,⁶ CHARLES H. NELSON,¹²
DOUGLAS RICHSTONE,¹³ DONNA WEISTROP,¹² AND BRUCE WOODGATE⁸

Received 2004 December 20; accepted 2005 May 9

ABSTRACT

We present *Hubble Space Telescope* (HST) spectroscopy of the nucleus of M31 obtained with the Space Telescope Imaging Spectrograph (STIS). Spectra that include the Ca II infrared triplet ($\lambda \simeq 8500 \text{ \AA}$) see only the red giant stars in the double brightness peaks P1 and P2. In contrast, spectra taken at $\lambda \simeq 3600\text{--}5100 \text{ \AA}$ are sensitive to the tiny blue nucleus embedded in P2, the lower surface brightness nucleus of the galaxy. P2 has a K-type spectrum, but we find that the blue nucleus has an A-type spectrum: it shows strong Balmer absorption lines. Hence, the blue nucleus is blue not because of AGN light but rather because it is dominated by hot stars. We show that the spectrum is well described by A0 giant stars, A0 dwarf stars, or a 200 Myr old, single-burst stellar population. White dwarfs, in contrast, cannot fit the blue nucleus spectrum. Given the small likelihood for stellar collisions, recent star formation appears to be the most plausible origin of the blue nucleus. In stellar population, size, and velocity dispersion, the blue nucleus is so different from P1 and P2 that we call it P3 and refer to the nucleus of M31 as triple.

Because P2 and P3 have very different spectra, we can make a clean decomposition of the red and blue stars and hence measure the light distribution and kinematics of each uncontaminated by the other. The line-of-sight velocity distributions of the red stars near P2 strengthen the support for Tremaine's eccentric disk model. Their wings indicate the presence of stars with velocities of up to 1000 km s^{-1} on the anti-P1 side of P2.

The kinematics of P3 are consistent with a circular stellar disk in Keplerian rotation around a supermassive black hole. If the P3 disk is perfectly thin, then the inclination angle $i \simeq 55^\circ$ is identical within the errors to the inclination of the eccentric disk models for P1+P2 by Peiris & Tremaine and by Salow & Statler. Both disks rotate in the same sense and are almost coplanar. The observed velocity dispersion of P3 is largely caused by blurred rotation and has a maximum value of $\sigma = 1183 \pm 201 \text{ km s}^{-1}$. This is much larger than the dispersion $\sigma \simeq 250 \text{ km s}^{-1}$ of the red stars along the same line of sight and is the largest integrated velocity dispersion observed in any galaxy. The rotation curve of P3 is symmetric around its center. It reaches an observed velocity of $V = 618 \pm 81 \text{ km s}^{-1}$ at radius $0''.05 = 0.19 \text{ pc}$, where the observed velocity dispersion is $\sigma = 674 \pm 95 \text{ km s}^{-1}$. The corresponding circular rotation velocity at this radius is $\sim 1700 \text{ km s}^{-1}$. We therefore confirm earlier suggestions that the central dark object interpreted as a supermassive black hole is located in P3.

Thin-disk and Schwarzschild models with intrinsic axial ratios $b/a \lesssim 0.26$ corresponding to inclinations between 55° and 58° match the P3 observations very well. Among these models, the best fit and the lowest black hole mass are obtained for a thin-disk model with $M_\bullet = 1.4 \times 10^8 M_\odot$. Allowing P3 to have some intrinsic thickness and considering possible systematic errors, the 1σ confidence range becomes $(1.1\text{--}2.3) \times 10^8 M_\odot$. The black hole mass determined from P3 is independent of but consistent with Peiris & Tremaine's mass estimate based on the eccentric disk model for P1+P2. It is ~ 2 times larger than the prediction by the correlation between M_\bullet and bulge velocity dispersion σ_{bulge} . Taken together with other reliable black hole mass determinations in nearby galaxies, notably the Milky Way and M32, this strengthens the evidence that the $M_\bullet\text{-}\sigma_{\text{bulge}}$ relation has significant intrinsic scatter, at least at low black hole masses.

We show that any dark star cluster alternative to a black hole must have a half-mass radius $\lesssim 0''.03 = 0.11 \text{ pc}$ in order to match the observations. Based on this, M31 becomes the third galaxy (after NGC 4258 and our Galaxy) in which clusters of brown dwarf stars or dead stars can be excluded on astrophysical grounds.

Subject headings: black hole physics — galaxies: individual (M31) — galaxies: nuclei

¹ Universitäts-Sternwarte, Scheinerstrasse 1, Munich D-81679, Germany; bender@usm.uni-muenchen.de, jthomas@usm.uni-muenchen.de.

² Max-Planck-Institut für Extraterrestrische Physik, Giessenbachstrasse, 85748 Garching-bei-München, Germany; bender@mpe.mpg.de.

³ Beatrice M. Tinsley Centennial Visiting Professor, University of Texas at Austin.

⁴ Department of Astronomy, University of Texas, RLM 15.308, C-1400, Austin, TX 78712; kormendy@astro.as.utexas.edu.

⁵ Computer Sciences Corporation, Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218; bower@stsci.edu.

⁶ National Optical Astronomy Observatory, P.O. Box 26732, Tucson, AZ 85726; green@noao.edu, lauer@noao.edu.

⁷ Emergent-IT, 1315 Peachtree Court, Bowie, MD 20721; danks@yahoo.com.

⁸ NASA Goddard Space Flight Center, Code 681, Greenbelt, MD 20771; gull@sea.gsfc.nasa.gov, woodgate@stars.gsfc.nasa.gov.

⁹ Herzberg Institute of Astrophysics, National Research Council of Canada, 5071 West Saanich Road, Victoria, BC V9E 2E7, Canada; john.hutchings@hia.nrc.ca.

¹⁰ Department of Physics and Astronomy, Rutgers University, P.O. Box 849, Piscataway, NJ 08855; cjoseph@physics.rutgers.edu.

¹¹ Department of Physics and Astronomy, Johns Hopkins University, Homewood Campus, Baltimore, MD 21218; kaiser@pha.jhu.edu.

¹² Department of Physics, University of Nevada, 4505 South Maryland Parkway, Las Vegas, NV 89154; cnelson@physics.unlv.edu, weistrop@physics.unlv.edu.

¹³ Department of Astronomy, University of Michigan, Dennison Building, Ann Arbor, MI 48109; dor@umich.edu.

1. INTRODUCTION

M31 was the second¹⁴ galaxy in which stellar dynamics revealed the presence of a supermassive black hole (BH; Kormendy 1987, 1988; Dressler & Richstone 1988). The spatial resolution of the discovery spectra was FWHM $\sim 1''$. Axisymmetric dynamical models implied BH masses of $M_{\bullet} = (1-10) \times 10^7 M_{\odot}$. The smallest masses were given by disk models, and the largest were given by spherical models.

In 1988, it was already known that axisymmetry is only an approximation to a more complicated structure. With Stratoscope II, Light et al. (1974) had observed that the nucleus is asymmetric. The brightest point is offset both from the center of the bulge (Nieto et al. 1986) and from the velocity dispersion peak (Dressler 1984; Dressler & Richstone 1988; Kormendy 1988). Then, using *Hubble Space Telescope* (HST), Lauer et al. (1993) discovered that the nucleus is double. The brighter nucleus, P1, is offset from the bulge center by $\sim 0''.5$. The fainter nucleus, P2, is approximately at the bulge center. Early concerns that an apparently double structure might only be due to dust were laid to rest when infrared images proved consistent with optical and ultraviolet images (Mould et al. 1989; Rich et al. 1996; Davidge et al. 1997; Corbin et al. 2001). These results were confirmed at higher resolution and signal-to-noise ratio (S/N) using WFPC2 (Lauer et al. 1998). With the discovery of the double nucleus, work on the central parts of M31 went into high gear.

Bacon et al. (1994, 2001) used integral-field spectroscopy to map the two-dimensional velocity field near the center of M31. They found that the kinematical major axis of the nucleus is not the same as the line that joins P1 and P2. The rotation curve is approximately symmetric about P2, i.e., about the center of the bulge. However, this is not the point of maximum dispersion. Instead, the brightest and hottest points are displaced from the rotation center by similar amounts in opposite directions.

The above results created two acute needs. First, the rich phenomenology of the double nucleus cried out for explanation. Second, the P1-P2 asymmetry raised doubts about BH mass measurements. This paper is mainly about the BH. HST allows us to take an important step inward by studying a blue cluster of stars embedded in P2. We introduce this cluster in § 1.1. Second, our spectroscopy of P1+P2 (§§ 2 and 3) provides further support for the preferred model of the double nucleus (Appendix). Since that model affects much of our discussion, we summarize it in § 1.2. For comprehensive reviews, see Peiris & Tremaine (2003) and Salow & Statler (2004).

1.1. P3: The Blue Star Cluster Embedded in P2

Nieto et al. (1986), using a photon-counting detector on the Canada-France-Hawaii Telescope (CFHT), were the first to illustrate that P2 is brighter than P1 at 3750 Å (contrast their Fig. 3 with Fig. 4 in Light et al. 1974; cf. Fig. 3 below). However, they did not realize this. Instead, they focused on the strong color gradient (bluer inward) and worried because this was inconsistent with published data. But these data were taken in the red or else had poor spatial resolution; they could not have seen the ultraviolet center. Nieto and collaborators found no problem with their data but concluded that “further observations are required to settle this question.”

King et al. (1992) confirmed the ultraviolet excess in the nucleus using the HST Faint Object Camera (FOC) at 1750 Å.

Using the same image, Crane et al. (1993a) illustrated that P2 is brighter than P1 but did not comment on this. Bertola et al. (1995) illustrated the same effect using FOC+F150W+F130LP images but again did not comment that it is P2, not P1, that is brighter in the ultraviolet.

Therefore, it was King et al. (1995) who discovered that P2 is much brighter than P1 in the ultraviolet. This result was again based on the 1750 Å FOC images. The blue light comes from a compact source that is embedded in P2 and that is similar in color and brightness to post-asymptotic giant branch (PAGB) stars seen elsewhere in the bulge (King et al. 1992; Bertola et al. 1995). King et al. (1995) proposed that the source might be nonthermal light from the weak active galactic nucleus (AGN) that is detected in the radio (Crane et al. 1992, 1993b), although they recognized that it could be a single PAGB star. Subsequently, Lauer et al. (1998) and Brown et al. (1998) resolved the source; its half-power radius is $\simeq 0''.06 = 0.2$ pc. Both papers argued that it is a cluster of stars. Lauer et al. (1998) combined the King et al. (1995) UV fluxes with optical fluxes to conclude that the source is consistent with an A star spectrum.

In this paper we present STIS spectra and show directly that the source is composed of A stars (§ 4). We also demonstrate that it is most consistent with a disk structure rather than with a dynamically hot cluster (§§ 5 and 6). Because the blue cluster is so distinct from P1+P2 in terms of stellar content and kinematics, we call it P3 and refer to the nucleus of M31 as triple.

The disk structure of P3 allows us to make a new and more reliable measurement of the central dark mass (§ 7). From the kinematics of P3 we also show that the dark object must be confined inside a radius $r \lesssim 0''.03 = 0.11$ pc. This implies that alternatives to a BH, such as a cluster of brown dwarf stars or stellar remnants, are inconsistent with the observations (§ 8).

1.2. The Eccentric Disk Model of P1+P2

Tremaine (1995) proposed what is now the standard model of P1 and P2. His motivation was the realization (see also Emsellem & Combes 1997) that the simplest alternative, an almost completed merger, is implausible. Two clusters in orbit around each other at a projected separation of $0''.49 = 1.8$ pc would merge in $\lesssim 10^8$ yr by dynamical friction. Instead, Tremaine proposed that both nuclei are parts of the same eccentric disk of stars. The brighter nucleus, P1, is farther from the BH and results from the lingering of stars near apocenter. The fainter nucleus, P2, is explained by increasing the disk density toward the center. A BH is required in P2 to make the potential almost Keplerian; only then might the alignment of orbits be maintained by the modest self-gravity of the disk.

Statler et al. (1999), Kormendy & Bender (1999, hereafter KB99), and Bacon et al. (2001) showed that the nucleus has the signature of the eccentric disk model. The most direct evidence is the asymmetry in $V(r)$ and $\sigma(r)$. Eccentric disk stars should linger at apocenter in P1; V and σ are observed to be relatively small there. The same stars should pass pericenter in P2, slightly on the anti-P1 side of the BH; the velocity amplitude is observed to be high on the anti-P1 side of the blue cluster. Because the PSF and the slit blur light from stars seen at different radii and viewing geometries, the apparent velocity dispersion should also have a sharp peak slightly on the anti-P1 side of the BH. All of the above papers demonstrated that the dispersion has a sharp peak in P2. KB99 showed further that the σ peak is slightly on the anti-P1 side of the blue cluster. Therefore, they suggested that the BH is in the blue cluster. Finally, KB99 demonstrated that the spectra and metal line strengths of P1 and P2 are similar to each other

¹⁴ The first was M32 (Tonry 1984, 1987). In retrospect, the resolution was barely good enough for a successful BH detection (Kormendy 2004); i.e., the BH was discovered essentially as early as possible.

TABLE 1
STIS INSTRUMENT CONFIGURATIONS

Parameter	Red Spectrum	Blue Spectrum
Detector gain ($e^- \text{ ADU}^{-1}$)	1.0	1.0
Grating	G750M	G430L
Wavelength range (\AA)	8272–8845	2900–5700
Reciprocal dispersion (\AA pixel^{-1})	0.56	2.73
Slit width (arcsec)	0.1	0.2
Comparison line FWHM (pixels)	3.1	3.5
$R = \lambda/\Delta\lambda$	4930	450
Instrumental dispersion σ_{instr} (km s^{-1})	56	284
Scale along slit (arcsec pixel^{-1})	0.051	0.051
Slit length (arcmin)	0.8	0.8
Integration time (s)	20790	2040

but different from those of the bulge. Therefore, P1 cannot be an accreted globular cluster or dwarf galaxy.

Peiris & Tremaine (2003) refined the eccentric disk model to optimize the fit to the higher resolution and more detailed ground-based spectroscopy now available. Even the Gauss-Hermite coefficients h_3 and h_4 , which were not used in constructing the model, were adequately well fitted. These models were then used to predict the kinematics that should be observed in our Ca triplet *HST* spectra of the red stars. This is a stringent test because the new models were used to predict observations taken at much higher resolution than those used to construct the models. Excellent fits were obtained. This is a resounding success of the eccentric disk model. The structural and velocity asymmetries of the nucleus can be explained almost perfectly if the eccentric disk is inclined with respect to the plane of the outer disk of M31. Here we publish the kinematic data used by Peiris & Tremaine (2003) in the above comparison (§ 3), and we revisit particularly interesting features of the STIS kinematics of P1+P2 in the Appendix.

The main shortcoming of the Peiris & Tremaine (2003) models is that they do not include the self-gravity of the stars in the eccentric disk. If the disk has a mass of 10% of the BH, then self-gravity is needed to keep the model aligned (Statler 1999). The most detailed such models are by Salow & Statler (2001, 2004). They model all available observations but do not fit the data as

well as the models by Peiris & Tremaine (2003). Other self-consistent models are based on N -body simulations (Bacon et al. 2001; Jacobs & Sellwood 2001); again, they reproduce only some of the observations. Sambhus & Sridhar (2002) use the Schwarzschild (1979) method to model the double nucleus. The above models differ in many details. For example, the Salow & Statler (2001, 2004) models precess rapidly, with pattern speeds of $36 \pm 4 \text{ km s}^{-1} \text{ pc}^{-1}$; the models of Sambhus & Sridhar (2002) precess at $16 \text{ km s}^{-1} \text{ pc}^{-1}$, and the simulations of Bacon et al. (2001) precess at only $3 \text{ km s}^{-1} \text{ pc}^{-1}$. Not surprisingly, the construction of dynamical models that include self-gravity is a challenge. The conclusion that such models are long lived is less secure than the result that they can instantaneously fit the photometry and kinematics of P1+P2. Tremaine (2001) gives a general discussion of slowly precessing eccentric disks.

Because of these complications, the BH mass in M31 has remained uncertain. Estimates of M_\bullet by Dressler & Richstone (1988), Kormendy (1988), Richstone et al. (1990), Bacon et al. (1994), Magorrian et al. (1998), KB99, Bacon et al. (2001), Peiris & Tremaine (2003), and Salow & Statler (2004) have ranged over a factor of about 3, $M_\bullet \simeq (3-10) \times 10^7 M_\odot$. These results are reviewed and error bars are tabulated on a uniform distance scale ($D = 0.76 \text{ Mpc}$) in Kormendy (2004). In this paper we show that an analysis of the UV-bright nucleus P3 allows us to estimate the BH mass independent of P1+P2.

2. STIS SPECTROSCOPY

The STIS CCD observations of M31 were obtained on 1999 July 23–24. The slit was aligned at P.A. = 39° . Other details of the STIS configuration are given in Table 1. We obtained a spectrum that includes the calcium triplet, $\lambda\lambda 8498, 8542,$ and 8662 , and one at $2700-5200 \text{ \AA}$ that includes several Balmer lines and Ca II H and K ($\lambda\lambda 3933$ and 3968). Both wavelength regions were observed because we wanted to analyze separately the double nucleus P1+P2 and the central blue cluster P3. Figure 1 shows that the double nucleus P1+P2 contributes almost all of the light at red wavelengths, while P3 dominates at 3000 \AA . The color difference between P1+P2 and P3 is illustrated further in the brightness cuts in Figure 3 below. The red spectrum was obtained using the $52'' \times 0''.1$ slit, while the blue spectrum was




FIG. 1.—STIS slit positions superimposed on the WFPC2 images from Lauer et al. (1998). The left panel shows the $0''.1$ slit position for the Ca II spectrum on the F555W image, and the right panel shows the $0''.2$ slit position for the blue spectrum on the F300W image. The images cover the central $6''.4 \times 6''.4$. North is $55''.7$ counterclockwise from up.

taken with the $52'' \times 0.2''$ slit. A wider slit was chosen for the blue spectrum to ensure that P3 would fall inside the slit. Figure 1 shows the placement of the slit relative to the WFPC2 F555W and F300W images from Lauer et al. (1998). These slit positions were determined by comparing the light profiles along the slit in our STIS spectra with brightness cuts through the WFPC2 images. We measured the slit positions to an accuracy of $0.005''$ for the red spectrum and $0.01''$ for the blue spectrum. The total integration time for the red spectrum was split into two exposures of approximately 1200 s each per *HST* orbit. M31's nucleus was shifted by 4.1 pixels along the slit between orbits.

The total integration time for the blue spectrum was split into three equal exposures within one *HST* orbit. The nucleus was shifted by 4.3 pixels between successive exposures. Wavelength calibrations (wavelcal) were interspersed among the galaxy exposures to allow wavelength calibration, including correction for thermal drifts. For the red spectrum, we obtained contemporaneous flat-field exposures through the same slit while M31 was occulted by the Earth. These provide proper calibration of internal fringing, which is significant at $\lambda \geq 7500 \text{ \AA}$ (see Goudfrooij et al. 1997).

The spectra were reduced as described in Bower et al. (2001). Unlike red spectra taken at $\lambda \geq 7500 \text{ \AA}$ with the G750M grating, blue spectra taken with the G430L grating are not affected by fringing. Consequently, we flat-fielded the G430L data using the library flat image from the STScI archive. The final reduced spectra have maximum S/N values of $S/N = 25$ (G750M) and 50 \AA^{-1} (G430L).

A stellar template spectrum is needed to measure the stellar kinematics implied by the galaxy spectra. For the red spectrum of M31, our template is the STIS spectrum of HR 7615 from Bower et al. (2001). They document the observational setup and data reduction for this spectrum. For the blue spectrum we used template A stars from Le Borgne et al. (2003), white dwarf stars observed in the Sloan Digital Sky Survey (SDSS; Kleinman et al. 2004) or modeled by Finley et al. (1997) and Koester et al. (2001), and spectral syntheses of various stellar population models by Bruzual & Charlot (2003). These sources were supplemented for checking purposes by using standard stars from Pickles (1998). Spectral resolution is not an issue for standard stars because the intrinsic width of the absorption lines in A-type stars is much larger than the instrumental width of STIS with the G430L grating and because the spectrum of P3 proves to have exceedingly broad lines.

3. KINEMATICS OF THE DOUBLE NUCLEUS P1+P2

The calcium triplet spectroscopy sees only the red giant stars that make up the double nucleus, P1+P2. It is blind to P3, which contributes essentially no light at $\lambda \simeq 8500 \text{ \AA}$. The kinematic properties of the red stars are illustrated in Figure 2.

In Figure 2, the spectrum of the bulge has been subtracted following procedures discussed in KB99. Bulge subtraction is analogous to sky subtraction in the sense that it removes the effects of a contaminating spectrum that is not of present interest. As shown in KB99, the bulge of M31 dominates the light distribution only at radii $r \geq 2''$. At $r < 1''$, it contributes about 20% of the light. So over the radii of interest in Figure 2, bulge stars are a minor foreground and background contaminant; they do not significantly participate in the dynamics of the double nucleus. It is routine to estimate the small contribution of bulge stars to the STIS red spectrum and to subtract it. Figure 2 is therefore a pure measure of the kinematics of the stars that make up the double nucleus.

Figure 2 also shows the bulge-subtracted nuclear kinematics measured with CFHT (KB99). Taking into account both the PSF




FIG. 2.—Panel 1 shows the double nucleus of M31 rotated $\sim 185^\circ$ clockwise with respect to Fig. 1. It is a $I+V+3000 \text{ \AA}$ composite from KB99. P1 is brighter than P2 in red light. Embedded in P2 is P3, i.e., a tiny cluster of blue stars that is invisible in I but brighter than P1 in the ultraviolet. The background image in panel 2 is a similar $V+3000 \text{ \AA}$ composite that better shows the small radius of P3. Panel 2 includes an I -band brightness cut along the P1-P2 axis (*lower line*) and a V -band cut through the blue cluster P3 (*upper line*). The filled circles are the brightness profile in the STIS spectrum; they are used to register the kinematics with the photometry in radius. Along the P1-P2 axis, radius $r = 0$ is chosen to be the center of P3 (note that in KB99 we centered the radius scale at $0.068''$, not P3). Panels 3 and 4 show velocity dispersions and rotation velocities along the P1-P2 axis after subtraction of the bulge. The ground-based points (*crosses*) are from the Subarcsecond Imaging Spectrograph (“SIS”) and the CFHT (KB99). The STIS data (*filled circles*) are Fourier quotient reductions. Bacon et al. (2001) made an independent reduction of our red STIS spectrum; it is consistent with ours.

and the slit, the effective Gaussian dispersion radius of the effective PSF was $\sigma_* = 0.297''$ (Kormendy 2004). The corresponding resolution of the STIS red spectroscopy is $\sigma_* = 0.052''$.

Confirming results of KB99, the dispersion profile of the red stars reaches a sharp peak slightly on the anti-P1 side of P3. The peak dispersion is higher at STIS resolution ($\sigma = 373 \pm 50 \text{ km s}^{-1}$) than at CFHT resolution ($\sigma = 287 \pm 9 \text{ km s}^{-1}$). The rotation curve is also asymmetric; the maximum rotation velocity is larger on the anti-P1 side than it is in P1. Again, the




FIG. 3.—Linear intensity cuts through the blue and red spectra of P1+P2+P3. Each cut is an average over the wavelength range given in the key. The contrast between the blue cluster P3 and the underlying red nucleus P2 is largest at 4000 Å. It is smaller at redder wavelengths because the stars in P3 are blue. It is smaller at bluer wavelengths because the spectrum of P3 has a strong Balmer break (Fig. 4). The two leftmost vertical dashed lines indicate the region in which the background spectrum was derived. The two rightmost vertical dashed lines indicate the radius range over which we averaged the background-subtracted P3 spectrum shown in Figs. 4, 5, and 6.

asymmetry is larger and the radius of maximum rotation is smaller at STIS resolution than at CFHT resolution. These observations are consistent with and provide further evidence for Tremaine’s (1995) model for the double nucleus as an eccentric disk of stars orbiting the central BH. The Appendix provides more detailed discussion.

4. THE INTEGRATED SPECTRUM OF P3

4.1. P3 Is Made of A-Type Stars

P3, the compact blue cluster, is illustrated in the two panels of images in Figure 2. It is embedded in P2 but is not concentric with it; the photocenter of P2 is $\sim 0''.03$ on the anti-P1 side of the blue cluster. The center of the bulge is slightly off in the opposite direction, i.e., toward P1 (see KB99 and discussion below). Note that we choose $r = 0$ to be the center of P3, whereas KB99 chose $r = 0$ to be the center of the bulge.

We obtained our STIS spectrum at $\lambda \simeq 3500\text{--}5000$ Å in part to study this issue. Over the above wavelength range, P3 provides a strong signal, much stronger than that indicated by the V -band brightness cut in Figure 2. Figure 3 shows brightness cuts through the red and blue STIS spectra in various wavelength ranges. The blue cluster is essentially invisible at 8300–8800 Å in the red spectrum. We assume that this spectrum provides the surface brightness profile of the underlying double nucleus. With respect to this profile, P3 is, in general, more prominent at bluer wavelengths.¹⁵ The contrast over P1+P2 is

¹⁵ P3 looks fainter at 4700–5100 Å in Fig. 3 than at 5500 Å (V -band) in Fig. 2. The reason is that Fig. 2 shows a brightness cut through the deconvolved V -band image from Lauer et al. (1998); this has higher spatial resolution than an un-deconvolved STIS spectrum. Also, the V -band cut is $0''.046$ wide, while the spectrum was obtained through a $0''.2$ wide slit.




FIG. 4.—Spectrum (*black line*) of the central $0''.2$ of the blue cluster P3. The superposed spectrum of the stars in the bulge and nucleus has been subtracted. Flux is in arbitrary linear units. In the bottom panel, the spectrum has been divided by a polynomial c_λ fitted to the continuum; it has been normalized to zero intensity and multiplied by the mean ratio γ of the line strength in the standard stars to that in P3. The colored lines show the spectra of an A0 dwarf star, an A0 giant star, and a Bruzual & Charlot (2003) starburst of age 200 Myr before (*top*) and after (*overplotted on the data*) broadening to the LOSVD that best fits the cluster spectrum. The fit was carried out with the Fourier correlation quotient program (Bender 1990).

highest at 3800–3950 Å. Then P3 gets less prominent at 3600–3750 Å; the reason turns out to be that the spectrum has a strong Balmer break (Fig. 4). The important conclusion from Figure 3 is that the spectrum of P3 is almost as bright as the underlying spectrum of P1+P2 at just the wavelengths where hydrogen Balmer lines are strongest.

It is therefore possible to extract a clean spectrum of P3 despite the short integration time and modest S/N. We averaged the spectrum of P3 over the $0''.2$ (i.e., four spectral rows) in which it is brightest (Fig. 3, *right pair of dashed lines*). We approximated the spectrum of the underlying P2 stars by averaging 14 rows of the spectrum on the anti-P1 side of P3 (Fig. 3, *left pair of dashed lines*). The 8300–8800 Å brightness cut was used to scale this average P2 spectrum to the P2 brightness underlying P3. The result was subtracted from the four-row average spectrum of P2+P3. The resulting spectrum of P3 is shown in black in Figures 4, 5, and 6.

The stellar population of P3 is dramatically different from that of P1 and P2. The spectrum in Figure 4 is dominated by Balmer absorption lines. At least five Balmer lines are visible, starting with H β at $\lambda_{\text{obs}} \simeq 4856$ Å. Also prominent is a strong Balmer break. In fact, the spectrum is very well matched by velocity-broadened spectra of A giant and dwarf stars. This confirms that the nucleus is made mostly of A-type stars as Lauer et al. (1998) and Brown et al. (1998) suggested.

4.2. The Remarkably High Velocity Dispersion of P3: The Supermassive Black Hole Is in the Blue Cluster

The blue cluster has a remarkably high velocity dispersion. Using an A0 dwarf star from Le Borgne et al. (2003) as a template,




FIG. 5.—This figure is analogous to Fig. 4, except that the spectrum of the blue cluster P3 is compared with Bruzual & Charlot (2003) starbursts of various ages given in the key. The fit to the red continuum is best for an age of ~ 510 Myr, but then the strengths of the Balmer lines H_n for $n \geq 5$ are wrong relative to the strengths of the redder lines. This problem gets worse for older starburst ages. Starbursts younger than 200 Myr are too blue; their Balmer breaks are too small to fit the observed spectrum.

the Fourier correlation quotient program (Bender 1990) gives a velocity dispersion of $\sigma = 962 \pm 105 \text{ km s}^{-1}$. An A0 giant star gives $\sigma = 984 \pm 107 \text{ km s}^{-1}$. A-type stars have intrinsically broad lines, but σ is so large that the difference between using giants and dwarfs is insignificant. The above fits are illustrated in Figure 4. The match to the lines and to the Balmer break is excellent. The results are robust; plausible changes in the intensity scaling of the P2 spectrum that was subtracted produce no significant change in σ .

The best-fitting 200 Myr old stellar population model (Fig. 5, § 4.3) gives a dispersion of $\sigma = 984 \pm 106 \text{ km s}^{-1}$. We adopt the average of the dispersion values given by the A dwarf star, the A giant star, and the 200 Myr old stellar population model; this gives $\sigma = 977 \pm 106 \text{ km s}^{-1}$ as our measure of the velocity dispersion of P3 integrated over the central $0''.02$.

Despite its tiny size (half-power radius $\simeq 0''.06 \simeq 0.2 \text{ pc}$; Lauer et al. 1998), P3 has the highest integrated velocity dispersion measured to date in any galaxy. The velocity dispersion of P3 is even larger than the line-of-sight velocity dispersion of the Sgr A* cluster in our Galaxy ($\sigma = 498 \pm 52$ to $840 \pm 104 \text{ km s}^{-1}$, depending on the sample of stars chosen; Schödel et al. 2003).¹⁶ The high velocity dispersion of P3 is especially remarkable in view of the observation (Fig. 2) that the velocity dispersion of the red stars *along the same line of sight* is only $\sim 250 \text{ km s}^{-1}$. The maximum velocity dispersion of P2, $373 \pm 48 \text{ km s}^{-1}$ at $\Delta r \simeq 0''.06$ on the anti-P1 side of the blue cluster, is much smaller than that of P3. Even the remarkably high velocity dispersion $\sigma = 440 \pm 70 \text{ km s}^{-1}$ measured in P2 by Statler et al. (1999) is much

¹⁶ Of course, the pericenter velocities of the innermost individual stars in our Galaxy are in some cases much larger. The current record is held by S0-16, which was moving at $12,000 \pm 2000 \text{ km s}^{-1}$ when it passed within 45 AU = 0.0002 pc = 600 Schwarzschild radii of the Galaxy's BH (Ghez et al. 2005).




FIG. 6.—Sample CMD of a 200 Myr old single-burst population with solar metallicity and a total luminosity of $M_V = -5.7$. The spectrum is dominated by stars of $\sim 10,000 \text{ K}$ temperature. The diagram has been generated using the synthetic CMD algorithm of the Instituto de Astrofísica de Canarias (Aparicio & Gallart 2004).

smaller than the velocity dispersion of P3. This confirms the conclusion of KB99 that the M31 supermassive BH is in the blue cluster.

4.3. Fit of a Starburst Spectrum to P3

The overall continuum slope of P3 is best fitted not by a single A-type star but rather by the spectrum (Bruzual & Charlot 2003) of a single starburst population (SSP in Figs. 4 and 5) of age $\sim 200 \pm 50 \text{ Myr}$ and solar metallicity. The blue continuum fit is essentially perfect; the red continuum fit is improved slightly over the single-star fits. Starburst spectra with a range of ages are shown in Figure 5. Using older starbursts allows us to fix the fit to the 5000 \AA continuum, but only at an unacceptable price: the bluest Balmer lines are no longer well fitted. Complicating the model further would be overinterpretation; the error in the red continuum fit could be due to imperfect P2 subtraction or to small amounts of dust. But it is clear that we cannot exclude some admixture of older stars. Reasonable changes in metallicity also do not affect the fit: metallicity changes are largely degenerate with age changes.

How many stars make up P3? For an absolute visual magnitude of $M_V \approx -5.7$ (Lauer et al. 1998), we estimate the answer in Figure 6, using IAC-STAR, the synthetic color-magnitude diagram (CMD) algorithm of the Instituto de Astrofísica de Canarias (Aparicio & Gallart 2004). A 200 Myr old, single-burst population of solar abundance implies that about 200 stars between spectral types A5 and B5 dominate the spectrum. The large number of stars at the same temperature of $\sim 10,000 \text{ K}$ explains why the spectrum of P3 is so similar to that of a single A0 star. Figure 6 also shows why P3 has a fairly smooth appearance, although surface brightness fluctuations are visible in Figure 8 below. Only a few red evolved stars are present, and they do not contribute significantly to the light of P3. Future observations with resolutions of about $0''.01$ should resolve the brightest stars close to the BH.

For a Salpeter (1955) initial mass function with a lower mass cutoff at $0.1 M_{\odot}$, the total number of stars on the main sequence at present is $\sim 15,000$; their total mass is about $4200 M_{\odot}$. If the burst originally produced stars up to $100 M_{\odot}$, then the initial total mass of P3 was $\sim 5200 M_{\odot}$. Given the inefficiency of star formation, the total gas mass required to form P3 probably was of the order $10^6 M_{\odot}$.

Forming stars so close to a BH is not trivial. It may be possible if $\sim 3 \times 10^6 M_{\odot}$ of gas could be concentrated into a thin disk of radius 0.3 pc and velocity dispersion 10 km s^{-1} . Then Toomre's (1964) stability parameter $Q \simeq 1$. It is not easy to see how such an extreme configuration could be set up, especially without forming stars already at larger radii. Well before the BH makes star formation difficult, the surface density of the dissipating and shrinking gas disk would get high enough so that the Schmidt (1959) law observed in nuclear starbursting disks (Kormendy & Kennicutt 2004, Fig. 21) would imply a very high star formation rate. This star formation would have to be quenched until the gas disk got small enough to form P3. And then the star formation would have to be very inefficient to put only $\sim 5200 M_{\odot}$ of the $\sim 3 \times 10^6 M_{\odot}$ of gas into stars. Similar considerations make it difficult to understand young stars near the Galactic center BH (e.g., Morris 1993; Genzel et al. 2003, Ghez et al. 2003, 2005). Nevertheless, young stars (or at least high-luminosity, hot stars) are present. Complicated processes of star formation (e.g., Sanders 1998) may not realistically be evaluated by a simple argument based on the Toomre Q instability parameter. So, if a dense enough and cold enough gas disk can be formed, star formation may be possible, even close to a supermassive BH.

4.4. Could the Hot Stars in P3 Result From Stellar Collisions?

The alternative to a starburst could be that the hot stars of P3 are formed via collisions between lower mass stars in P3 or even in P1+P2. Yu (2003) argues that the collision timescales are too long to be of interest. It would be interesting to revisit this issue given the conclusion of § 6.1 that P3 is a cold stellar disk. In any case, it is worth noting that the conversion of, say, a high-mass, $0.5 M_{\odot}$ main-sequence star in P2 into an A star requires merging roughly six stars without mass loss. It is not easy to see how the A stars in P3 could originate by collisions.

Thus, the situation in P3 is similar to that in our Galaxy. No explanation of the hot stars looks especially plausible.

4.5. P3 Is Not Made of White Dwarf Stars

Finally, we need a sanity check to make sure that we are not completely misinterpreting the observations. Dynamically, we detect a $10^8 M_{\odot}$ central dark object. This is associated with a tiny and faint nucleus comprised of hot stars that have extraordinarily broad absorption lines. White dwarf stars have extraordinarily broad absorption lines. If they are not too old, they can easily have an A-type spectrum, and if they are not too young, they can easily contribute mass without contributing much light. It is natural to wonder, could P3 be a cluster of white dwarfs? Could they simultaneously explain the broad-lined, A-type spectrum and the central dark mass? This possibility is not excluded by stellar collision or cluster evaporation timescales (Maoz 1995, 1998).

Figures 7 and 8 show that P3 cannot be made of white dwarfs. Figure 7 compares the spectrum of P3 with that of a typical DA white dwarf observed in the SDSS. The star was chosen to have Balmer line strengths comparable to those in P3. It is approximately the best match to P3 that can be achieved with white dwarf spectra. Its lines are narrower than those of P3, so we can fit the observed line widths (Fig. 7, bottom panel) with




FIG. 7.—Spectrum of P3 fitted with approximately matched observed and model white dwarf spectra. The observed spectrum is from the SDSS (Kleinman et al. 2004; see <http://www.sdss.org>). The model spectra used in Figs. 7 and 8 are from Finley et al. (1997) and Koester et al. (2001). The fits of white dwarf spectra to P3 are significantly worse than the ones in Figs. 4 and 5. The absorption lines of the white dwarfs are intrinsically too strong, and the white dwarfs fail completely to fit the large Balmer continuum break in the P3 spectrum. However, the implied velocity dispersion, $\sigma = 885 \pm 126 \text{ km s}^{-1}$, is consistent with our adopted value, $\sigma = 977 \pm 106 \text{ km s}^{-1}$.

$\sigma = 885 \pm 126 \text{ km s}^{-1}$. That is, this relatively narrow-lined white dwarf gives a dispersion similar to those implied by main-sequence and giant A stars. The fit to the line widths is less good than the fit provided by A0 V stars, but it is not inconsistent with our low-S/N spectrum of P3. If we had only the spectrum of this white dwarf as observed over the relatively narrow wavelength region redward of the Balmer break, we could not exclude white dwarf stars.

However, the continua of white dwarf stars do not fit the large Balmer break in P3. SDSS J094624.30+581445.4 (Fig. 7) does not show this: it and most other white dwarfs have not been observed at blue enough wavelengths to reach the Balmer break. Therefore, we resort to model spectra kindly provided by Detlev Koester (Finley et al. 1997; Koester et al. 2001). Figure 7 shows a model spectrum that has line profiles similar to those in the observed white dwarf. The price of having narrow enough lines to fit the absorption lines in P3 is that there is essentially no Balmer break. Such a star cannot fit the continuum of P3. This result is very robust; it is not affected by uncertainties in the subtraction of the spectra of P1+P2.

Choosing different white dwarf parameters does not solve this problem. No combination of temperature and gravity allows a simultaneous match to the Balmer line strengths and the Balmer break. Figure 8 shows fits of model white dwarf spectra with temperatures $T = 7000, 8000, 10,000,$ and $12,000 \text{ K}$. For each temperature, we try surface gravities of $10^7, 10^8,$ and 10^9 cm s^{-2} .

Temperature $T \simeq 7000 \text{ K}$ is too cold. The stellar lines are too weak. Not surprisingly, these stars have no Balmer break at all. Despite the bad continuum fit, the narrow lines in the white dwarf templates give dispersions $\sigma = 945 \pm 103, 987 \pm 107,$ and $1063 \pm 115 \text{ km s}^{-1}$ that are consistent with our adopted result.




FIG. 8.—Fits to the spectrum of P3 (black line) of model white dwarf spectra (red line) with temperature $T_{\text{eff}} = 7000, 8000, 10,000,$ and $12,000$ K (see the key). At each temperature, surface gravities of $g = 10^7, 10^8,$ and 10^9 cm s^{-2} are used. The green line shows the fit of an A0 dwarf star. Compare Fig. 6.

At $T = 8000$ K, the fit to the lines is better, although not as good as for A0 dwarf or giant stars. The dispersion remains high ($\sigma = 930 \pm 101, 929 \pm 101,$ and 952 ± 103 km s^{-1}). Again, the Balmer break in the white dwarfs is too weak.

At $T = 10,000$ K, the stellar lines are much broader. The fit to P3 is acceptable after scaling the line strengths. For $\log g = 7, 8,$ and 9 , $\sigma = 784 \pm 120, 769 \pm 134,$ and 821 ± 150 km s^{-1} , respectively. Note that without line-strength scaling, the broadened white dwarf spectrum does not fit the galaxy. And, even though the lines are now strong enough when $\log g$ is large to produce a Balmer break, it is still too small to fit the spectrum of P3. The green line emphasizes how much

an A0 V star fits the spectrum of P3 better than does any white dwarf.

Increasing the temperature further is counterproductive. At $T = 12,000$ K, the lines are too strong and too broad to fit P3, although we still obtain high dispersions ($\sigma = 705 \pm 144, 676 \pm 166,$ and 680 ± 190 km s^{-1}). Even high temperatures do not produce strong enough Balmer breaks.

We conclude that no spectral synthesis of white dwarf stars of different temperatures or gravities would fit P3. The ones that fail least badly, those that fit the lines but not the Balmer break, imply velocity dispersions that are consistent with values derived from A0 dwarf or giant stars.




FIG. 9.—*Left to right*: F300W image of the blue nucleus, P3, superposed on nuclei P1 and P2; F300W image of P3 after subtraction of the F555W image intensity scaled to the F300W image in the first panel; PSF-convolved inclined disk model for P3; and difference between images in the second and third panels showing the quality of the model and the residual surface brightness fluctuations. All images are $2''.5 \times 2''.5$. North is $55^\circ.7$ counterclockwise from up, as in Fig. 1.

For a compact cluster of white dwarfs to be a viable alternative to a supermassive BH, it must be dark. That is, it must be old. We explore this option further in § 8.

5. LIGHT DISTRIBUTION OF P3

For a dynamical analysis of P3 (§ 6), we need its light distribution with P1+P2 subtracted. To derive this, we scaled the *HST*

F555W image to the *HST* F300W image such that P1 disappeared after subtraction. The resulting image of P3 is shown in Figure 9. We then fitted P3 with Sérsic (1968) models,

$$I(r) = I_0 \exp\left[-(r/r_0)^{1/n}\right],$$

convolved with the *HST* point-spread function (PSF) as in Lauer et al. (1998). The PSF was constructed from two exposures of the standard star GRW +70D5824 (u2tx010at, u2tx020at). The free parameters in the fit were central surface brightness SB_0 , scale length $r_0 = (a_0 b_0)^{1/2}$ (where a and b are semimajor and semiminor axis, respectively), Sérsic n , position angle P.A., ellipticity $1 - b/a$, and center coordinates. Individual faint pointlike sources in the outskirts of P3 were masked before fitting. The best fit over the radius range $r < 0''.3$ was obtained for Sérsic index $n = 1$, major-axis scale length $a_0 = 0''.1 \pm 0''.01$, PSF-convolved ellipticity $1 - b/a = 0.33 \pm 0.03$, and position angle P.A. = $63^\circ \pm 2^\circ$ (this is 119° counterclockwise from vertical in Fig. 9).

The PSF-convolved model and the difference between P3 and the model are illustrated in Figure 9. We also compare the model and P3 with respect to their isophotal parameters. Figure 10 shows the surface brightness, ellipticity, and position angle profiles of the observed P3, the model, and the PSF-convolved model. Also shown are deconvolved P3 profiles, which were obtained from 15 iterations with the Richardson-Lucy deconvolution algorithm implemented in the ESO MIDAS package. The deconvolved surface brightness profile obtained here agrees well in shape with the one by Lauer et al. (1998). Figure 10 shows that the model represents P3 reasonably well, especially over the radius range for which we can analyze the kinematics (§ 6). Surface brightness fluctuations become large at radii beyond $0''.25$. Still, the model is adequate out to $\sim 0''.4$.

If P3 is an inclined, thin disk, then the observed ellipticity implies an inclination $i = 55^\circ \pm 2^\circ$. This is compatible with the




FIG. 10.—Observed radial profiles (*red line*) of P3 surface brightness SB, ellipticity $1 - b/a$, and position angle P.A. vs. semimajor axis a . Lucy-deconvolved profiles are shown in green. The *HST* PSF is shown in light blue (with arbitrary zero point). The inclined disk model before and after convolution with the *HST* PSF is represented by blue and black lines, respectively. The observed profiles are oversampled: neighboring points are not independent.

TABLE 2
PARAMETERS OF THE THIN-DISK MODEL OF P3

Parameter	Value
m_{F300W}	18.6 ± 0.1
M_{F300W}	-5.8 ± 0.1
Sérsic n	1
Scale length a_0 (arcsec)	0.1 ± 0.01
$SB_{0, F300W}$ (face-on) (mag arcsec $^{-2}$)	15.6 ± 0.1
Inclination (deg)	55 ± 2
Position angle (deg)	63 ± 2

inclination of the eccentric disk P1+P2: Peiris & Tremaine (2003) derive $i = 54^\circ$, and Bacon et al. (2001) get $i = 55^\circ$. The model parameters of P3 are summarized in Table 2. Whether P3 really is a thin disk can only be checked with kinematical data. We discuss these in the next section.

6. DYNAMICS OF P3

Figure 11 shows the rotation velocity and velocity dispersion profiles of P3. Table 3 lists the data. We used FCQ for the analysis but did not fit the h_3 and h_4 Gauss-Hermite parameters because the S/N of the data is only $\sim 3 \text{ \AA}^{-1}$. Outside of the central pixel, P3 rotates rapidly, with an observed amplitude of $573 \pm 61 \text{ km s}^{-1}$ (weighted mean of all points with $|r| > 0''.01$). P3 rotates in the same sense as P1+P2. The apparent velocity dispersion drops from $\sim 1200 \text{ km s}^{-1}$ in the central pixel to $< 500 \text{ km s}^{-1}$ at $r = 0''.15 = 0.55 \text{ pc}$. These values are consistent with the velocities seen in the extreme wings of the line-of-sight velocity distribution (LOSVD) of the red stars at $r \sim -0''.1$ (see the Appendix). The kinematic data securely locate the BH at the center of P3 with an uncertainty of about $\frac{1}{3}$ of a pixel, i.e., $0''.02 = 0.07 \text{ pc}$.

We wish to combine the surface brightness data (Table 2) and the kinematic data (Table 3) to make dynamical models. Because the pixel size, slit width, and PSF are all similar to the size of P3, unresolved rotation must contribute to the apparent velocity dispersion. Actually, almost all of the light of P3 falls into the slit. Despite this and despite the modest apparent flattening, P3's apparent rotation velocity and velocity dispersion are similar. Therefore, it is reasonable to expect that P3 is an intrinsically flat object with $V \gg \sigma$.

For these reasons, we first model P3 as a flat disk with an exponential profile and an inclination $i \sim 55^\circ$ (§ 6.1). Then (§ 6.2), we explore more nearly edge-on models in which P3 has some intrinsic thickness.

6.1. P3 as a Flat Exponential Disk

We construct a dynamical model in which we assume that P3 is a flat disk with the parameters in Table 2 and a negligible intrinsic velocity dispersion. The BH affects the structure of the galaxy interior to $r_{\text{cusp}} \simeq GM_\bullet/\sigma^2 = 5''.6[M_\bullet/(10^8 M_\odot)]$, where G is the gravitational constant and $\sigma = 145 \text{ km s}^{-1}$ (Kormendy 1988) is the velocity dispersion of the bulge just outside the region affected by the BH. Since P3 is tiny compared to r_{cusp} , the BH dominates the gravitational potential. The distribution of the stars is completely constrained by the photometry, so the only free parameter is the BH mass. To compare the model with the observed rotation and velocity dispersion profiles, we convolve the Keplerian velocity field with the PSF and integrate it over the $0''.2$ slit width and $0''.05$ CCD pixels (see Fig. 11, *top middle panel*). This is done with small subpixels to obtain smooth profiles of rotation velocity and velocity dispersion.

Figure 11 shows the results. The observed rotation and dispersion profiles (*open and filled symbols*) are well matched by the model (*dotted lines*). Estimating the mass of the BH is straightforward because M_\bullet is the only free parameter. The best fit gives $M_\bullet = (1.4 \pm 0.2) \times 10^8 M_\odot$. The reduced χ_n^2 is ~ 1 (Fig. 12).

The BH mass derived with the thin-disk model does not depend significantly on inclination over the range allowed by the photometry ($\pm 2^\circ$). Changing the inclination away from the best value increases V and decreases σ or vice versa. Then χ^2 increases slightly, but the shape of the χ^2 distribution as a function of BH mass does not change significantly. We also varied the

scale length of the P3 disk, its total luminosity, and its position angle on the sky within the errors. There was no significant effect on M_\bullet . The total luminosity and mass of P3 are irrelevant provided that the BH dominates the potential. The position angle would have to be changed well beyond its estimated errors to achieve a visible effect on the velocities. Changing the radial scale length redistributes light and makes the rotation and dispersion profiles flatter or steeper. Within the errors, M_\bullet is not affected.

The circular velocities for the thin-disk model are shown in the top panel of Figure 11. Future observations that resolve individual stars should see velocities of $1000\text{--}2000 \text{ km s}^{-1}$. Such measurements can also test how much the observed velocities and dispersions are affected by shot noise due to the small number of stars in P3. Checking how close to circular the disk really is will also be important.

If P3 is a thin stellar disk, can it be stable? The answer is yes, as long as its stellar mass is not very much larger than $5200 M_\odot$. Even relatively small dispersions will not lead to significant two-body relaxation. Using equation (8-71) in Binney & Tremaine (1987), we obtain relaxation times of the order of a Hubble time. Moreover, the critical velocity dispersion for local stability (Toomre 1964) is small, $\sigma_{\text{crit}} \ll 1 \text{ km s}^{-1}$. This is a consequence of the fact that the potential is dominated by the BH. That is, the P3 disk is dynamically analogous to Saturn's rings rather than to a self-gravitating disk. Therefore, if earlier starbursts contributed mass without affecting its present spectrum, the P3 stellar disk is likely to be locally stable and immune from two-body relaxation. And if P3 consists only of young stars, then it has not had time for significant dynamical evolution.

6.2. P3 Schwarzschild Models

To investigate the effect on M_\bullet of allowing P3 to have some thickness in the axial direction z and therefore to be more nearly edge-on than $i = 55^\circ$, we fitted Schwarzschild (1979) models to the photometric and kinematic data. We used the regularized maximum entropy method as implemented by Gebhardt et al. (2000a, 2003) and Thomas et al. (2004). The program was constrained to reproduce the observed surface brightness distribution of P3. We considered three inclinations $i = 58^\circ, 66^\circ$, and 90° , corresponding to intrinsic axial ratios of P3 of 0.26, 0.44, and 0.57, respectively. BH masses were varied until the kinematic data were reproduced as well as possible, as indicated by the χ^2 values in Figure 12.

In the Schwarzschild code, phase space is quantized on a polar grid that is not optimized for closed orbits. It is therefore helpful if the orbits are not quite closed. For this reason, we did not use a point mass for the central dark object but rather used a Plummer sphere with a half-mass radius $r_h = 0''.01$. Given the spatial resolution of the data, this is essentially equivalent to a BH (see Fig. 14 below).

Models that put significant weight on entropy maximization did not fit the kinematics. They rotated too slowly because they contained retrograde orbits. This is expected because entropy maximization is not appropriate for highly flattened systems with strong rotational support.

Switching off the entropy maximization (this corresponds to a high regularization parameter in Thomas et al. 2004) results in better fits. Figure 12 shows χ^2 values as a function of inclination and dark mass M_\bullet . We conclude that the lowest inclination, $i = 58^\circ$, is preferred by $\Delta\chi^2 \approx 2$ relative to the $i = 66^\circ$ model and with higher significance relative to the more inclined models.




FIG. 11.—The three color images at the top show the PSF-broadened thin-disk model of P3. The images cover $0''.5 \times 0''.5$. Shown from left to right are (1) P3 surface brightness (intensities range from 0 [black] to 1 [white]); (2) P3 rotation velocity field with the slit and radial bins superposed (the velocity amplitudes range from -700 [black] to $+700$ km s⁻¹ [white]); and (3) P3 velocity dispersion, ranging from 150 [black] to 1000 km s⁻¹ [white]. The panels of plotted data points show the P3 radial profiles (red) of rotation velocity (bottom) and velocity dispersion (middle), folded around P3's center. Open and filled symbols are from opposite sides of the center. The sense of rotation is the same as for the eccentric disk P1+P2. The top panel shows the best-fitting Keplerian circular velocity curve as a dashed line. It implies a BH mass of $\sim 1.4 \times 10^8 M_\odot$. Convolution of the circular velocity field with the PSF and integrating it over the pixel size and slit width yields the model rotation and dispersion profiles shown as dotted lines in the bottom and middle panels.

TABLE 3
 KINEMATICS OF P3

Radius (arcsec)	V (km s ⁻¹)	ΔV (km s ⁻¹)	σ (km s ⁻¹)	$\Delta\sigma$ (km s ⁻¹)
-0.16	525	197	237	233
-0.11	616	144	337	170
-0.06	582	111	583	131
-0.01	170	169	1183	200
+0.04	-659	117	777	139
+0.09	-387	179	769	211
+0.14	-420	273	505	322

Rotation velocity and velocity dispersion profiles for the lowest χ^2 model at each inclination are shown in Figure 13. Reassuringly, the $i = 58^\circ$ Schwarzschild model most nearly resembles the $i = 55^\circ$ thin-disk model. The fits then become progressively more different (and less good) as the models are made more edge-on.

Higher inclinations require higher BH masses. The reason is that, at higher inclinations, line-of-sight integration through the nearly edge-on, thick disk includes stars at relatively large radii that move mostly across, not along, the line of sight. They reduce the velocity moments and consequently require higher M_\bullet to match the observed rotation velocities. The preferred BH mass for the $i = 58^\circ$ and 66° Schwarzschild models is $\sim 2 \times 10^8 M_\odot$. The highest BH mass that is consistent with the data to within $\sim 1 \sigma$ is given by the $i = 58^\circ$ Schwarzschild model and is $\sim 2.3 \times 10^8 M_\odot$ (Fig. 12). The lowest BH mass implied by the dynamics of P3 is given by the thin-disk, $i = 55^\circ$ model and is $\sim 1.2 \times 10^8 M_\odot$ (§ 6.1).

It is instructive to examine the $i = 58^\circ$ Schwarzschild model in more detail. Figure 14 shows its velocity moments. Rotation dominates the dynamics; V_ϕ is $\lesssim 20\%$ smaller than the circular velocity. At radii $r \gtrsim 0.1$, the model is approximately isotropic.




FIG. 12.—The χ^2 profiles for models of P3. The black lines refer to the thin-disk model with inclination 55° , while the colored lines show three Schwarzschild models with inclinations 58° , 66° , and 90° . The dashed lines assume that the MDO is a Plummer sphere with radius $0.01''$; the solid black line assumes a BH.




FIG. 13.—Rotation velocities and velocity dispersions of P3 as in Fig. 11 (red open and filled symbols). Overplotted as colored dashed lines are three Schwarzschild models with different inclinations and BH masses. The thin-disk model of Fig. 11 is shown as a dotted black line.

To provide the thickness that is necessitated by the inclination $i > 55^\circ$, σ_z then increases substantially inward, although it remains smaller than the rotation velocity. The difference between the adopted Plummer model and a central point mass is small except for the innermost data point.

Figure 15 shows the corresponding orbit structure. As expected for a Schwarzschild model that is not too different from the thin-disk model, retrograde orbits are strongly suppressed. However, as indicated by Figure 14, noncircular orbits get significant weight in order to produce an axial ratio $b/a \simeq 0.26$. As expected, this happens more near the center than at larger radii. However, nearly circular orbits dominate; otherwise, V_ϕ would not be almost equal to V_c in Figure 14.




FIG. 14.—Major-axis velocity moments of the best $i = 58^\circ$ Schwarzschild model for P3, corresponding to $M_\bullet = 2 \times 10^8 M_\odot$. The MDO is a Plummer sphere with half-mass radius $r_h = 0.01''$; its circular velocity is shown as a solid black line. A BH of the same mass would produce the Keplerian circular velocities indicated by the dashed curve.




FIG. 15.—Orbit structure of the Schwarzschild model with $i = 58^\circ$, Plummer model dark mass $M_\bullet = 2 \times 10^8 M_\odot$, and half-mass radius $r_h = 0''.01$. For each orbit, the orbit weight w_i per phase space volume is shown as a function of the z -component of its angular momentum L_z normalized by the angular momentum L_c of the circular orbit that has the same energy. In this figure only, r is the average of the pericenter and apocenter radii of the orbit. Note that at all radii, only prograde orbits are significantly populated.

6.3. Summary: Comparison of P3 and P1+P2

We conclude that the triple nucleus of M31 is made of two nested, disklike systems. The P1+P2 disk is elliptical, has a radius of about 8 pc, and consists of old, metal-rich stars. If it is thin, it has an inclination of $\sim 54^\circ$ and a major-axis position angle of $\sim 43^\circ$ (Peiris & Tremaine 2003). The P3 disk is approximately circular and has a radius of about 0.8 pc. If it is thin, P3 has an inclination $i \sim 55^\circ$ that is the same as that of P1+P2. P3's major-axis position angle at $r \leq 0''.25$ is $\sim 63^\circ \pm 2^\circ$. That is, the inner P3 disk is slightly tilted with respect to the P1+P2 disk but is relatively close to the kinematical major axis, P.A. $\sim 56^\circ$, found by Bacon et al. (2001). At $r > 0''.25$, the major axis of P3 twists to $\sim 40^\circ$, essentially the position angle of P1+P2. The nested disks rotate in the same sense and have almost parallel angular momentum vectors.

7. THE MASS OF THE CENTRAL DARK OBJECT

We have demonstrated that disklike models for P3 fit both the photometry and the kinematics of P3 exceedingly well. This allowed an estimate of M_\bullet that is independent of all previous determinations. Besides BH mass, only inclination is a free parameter in the fit to the rotation curve and the dispersion profile (Figs. 11 and 13).

Could systematic effects cause additional errors that are not included in the statistical errors, especially toward low BH masses? We mentioned in the previous section that some clumpiness in the distribution of stars could be hidden by PSF blurring and may affect the measured velocities and dispersions. We carried out a simple check for this effect by fitting subsamples of data points. For example, a fit to just the innermost three points in Figure 11 typically yields BH masses about 15% higher, while omitting these three points results in 25% lower masses. All values obtained in this way fell in the range allowed by the χ^2 profiles in Figure 12, so this effect does not seem to be very important.

Some noncircularity of the P3 disk could be hidden as well. P3 could contain stars on elongated orbits that have their pericenters within the range of the kinematic data ($0''.15$) but apocenters spread out over radii well beyond this radius. Figure 9 shows faint blue stars that could be such objects. This would imply that P3's velocity amplitudes are increased by rotation velocities that are faster than circular. It is difficult to estimate the size of this effect, but pericenter velocities of very radial stars can be at most a factor of 2 larger than pericenter velocities of stars on nearly circular orbits. Averaging over a set of orbits will reduce this number considerably. And if we wanted to fully exploit this effect, many more stars of P3 would have to be found outside of $\sim 0''.18$ than inside, which is in contradiction with the observations. The Schwarzschild models also show that this trick does not work well. More radially biased models (obtained with higher entropy weighting and not shown here) require larger BH masses. Finally, if P3 originated in a star-forming gas disk, it could not contain nearly radial orbits, and we noted above that subsequent internal evolution of the P3 disk should be slow.

So, very special circumstances would be required to decrease M_\bullet below $1 \times 10^8 M_\odot$. On the high-mass end, the BH mass grows with increasing inclination of the model. However, the χ^2 values become unacceptably large for inclinations above $\sim 60^\circ$; therefore, it is unlikely that the BH mass is significantly larger than $\sim 3 \times 10^8 M_\odot$. Viable models for P3 are found in the inclination range $55^\circ < i < 58^\circ$ and in the BH mass range $1.1 \times 10^8 M_\odot < M_\bullet < 2.3 \times 10^8 M_\odot$. The upper limit takes into account that the Schwarzschild models were calculated assuming a massive dark object (MDO) with $r_h = 0''.01$ and not a BH; the upper limit for a BH is $\approx 0.2 \times 10^8 M_\odot$ lower than for an MDO with $r_h = 0''.01$. The best-fit and at the same time lowest BH mass of $M_\bullet = 1.4 \times 10^8 M_\odot$ is obtained for the thin-disk model. This model is also preferred on astrophysical grounds, if P3 formed out of a thin gaseous disk. Therefore, our best estimate for the mass of the supermassive BH in M31 is $M_\bullet = 1.4^{+0.9}_{-0.3} \times 10^8 M_\odot$.

How does this compare with previous results? The BH mass has now been estimated by five largely independent techniques: (1) standard dynamical modeling that ignores asymmetries, (2) the KB99 center-of-mass argument that depends on the asymmetry of P1+P2, (3) the Peiris & Tremaine (2003) nuclear disk model that explains the asymmetry of P1+P2, (4) full dynamical modeling that takes into account the self-gravity of the P1+P2 disk (Salow & Statler 2004), and (5) dynamical modeling of the blue nucleus P3, which is independent of P1+P2. The good news is that all methods require a dark mass with high significance. The bad news is that some of the results differ by more than two standard errors. In particular, the disagreement between the KB99 center-of-mass argument and the P3 models presented here is a concern. We therefore revisit the KB99 derivation in the subsection below. The models that best fit both the photometry and the kinematics, the Peiris & Tremaine (2003) eccentric disk model of P1+P2 and our thin-disk model of P3, agree within the errors and favor a high BH mass of $M_\bullet \sim 1 \times 10^8 M_\odot$. We also note that a higher BH mass can be accommodated more easily in almost all models than a lower BH mass.

The mass of the M31 BH derived here is a factor of ~ 2.5 above the ridgeline of the correlation between M_\bullet and bulge velocity dispersion σ_{bulge} (Ferrarese & Merritt 2000; Gebhardt et al. 2000b). Using the Tremaine et al. (2002) derivation,

$$\log\left(\frac{M_\bullet}{M_\odot}\right) = 8.13 + 4.02 \log\left(\frac{\sigma_{\text{bulge}}}{200 \text{ km s}^{-1}}\right),$$

$\sigma_{\text{bulge}} \simeq 160 \text{ km s}^{-1}$ implies that $M_{\bullet} \simeq 5.5 \times 10^7 M_{\odot}$. We derive $M_{\bullet} = 1.4^{+0.9}_{-0.3} \times 10^8 M_{\odot}$. Tremaine et al. (2002) already found significant scatter in the M_{\bullet} - σ_{bulge} relation at low masses. With the increased BH mass for M31, scatter has become even more prominent. Considering, in addition to M31, only the two closest other supermassive BHs, i.e., M32 and the Galaxy, we get the following: M32 has $\sigma_{\text{bulge}} \sim 75 \text{ km s}^{-1}$, a predicted $M_{\bullet} = 2.6 \times 10^6 M_{\odot}$, and an observed $M_{\bullet} = (2.9 \pm 0.6) \times 10^6 M_{\odot}$ (Verolme et al. 2002; corrected to distance 0.81 Mpc from Tonry et al. 2001). Our Galaxy has $\sigma_{\text{bulge}} \sim 103 \text{ km s}^{-1}$, a predicted $M_{\bullet} = 9.4 \times 10^6 M_{\odot}$, and an observed $M_{\bullet} = (3.7 \pm 0.2) \times 10^6 M_{\odot}$ (Ghez et al. 2005). So M31, M32, and our Galaxy have BH masses that are 2.5 times larger than, consistent with, and 3 times smaller than the ridgeline of the M_{\bullet} - σ_{bulge} relation, respectively. This is strong indication for significant intrinsic scatter in the M_{\bullet} - σ_{bulge} relation, at least at the low-mass end.

7.1. Black Hole Mass from the Center-of-Mass Argument: KB99 Revisited

KB99 estimated that the center of P3 (Fig. 16, *blue filled circle*) is offset from the bulge center (*horizontal dashed line*) by about $0''.06$. They then assumed that the central dark object is in P3 and estimated its mass based on the assumption that the combined system, BH+P1+P2+P3, is in dynamical equilibrium. That is, they assumed that the center of mass of BH+P1+P2+P3 is at the center of the bulge. Then M_{\bullet} is inversely proportional to its distance from the bulge center and, if the mass of P3 is negligible, is proportional to the mass of P1+P2. The latter was given by the light distribution and the measured mass-to-light ratio $M/L_V = 5.7$. The resulting BH mass was $M_{\bullet} \simeq (3.3 \pm 1.5) \times 10^7 M_{\odot}$. This is at the low end of the range of published values and a factor of ~ 4 smaller than the value derived here. As the stellar M/L_V can hardly be a factor of 4 larger, two explanations are possible for the discrepant BH masses: (1) P3 and the BH are a factor of 4 closer to the bulge center than KB99 derived, or (2) the BH+P1+P2+P3 system is not in equilibrium with respect to the bulge center.

We believe that the observations can be consistent with equilibrium and that the distance of P3 to the bulge center was overestimated by KB99 for the following reasons.

Figure 16 revisits the center-of-mass argument. It is reproduced from KB99 and shows their estimate of the position of the center of the bulge as the dashed line at $X = 0$. The isophote center coordinate X is measured along the line joining P1 and P2. A conclusion about the position of the center of the bulge depends on the radius range chosen in which to average isophote X -values. KB99 calculated the average at $2''.9 < a < 25''.0$. A larger radius range was not possible because of the small size of the high-resolution images used. If $M_{\bullet} \sim 3 \times 10^7 M_{\odot}$, then the above radius range is no problem: it is beyond the radii affected by the BH. However, if M_{\bullet} is as big as $1.4 \times 10^8 M_{\odot}$, then $r_{\text{cusp}} \simeq 7''.2$ and it is necessary to calculate the mean bulge X at larger radii.¹⁷

If we calculate the bulge center outside of $r_{\text{cusp}} \simeq 7''.2$, we obtain a mean X -position of $-0''.033$ in Figure 16, i.e., halfway between the KB99 bulge center and P3. Note that, unlike KB99, we do not limit the averaging to points with $a < 25''.0$ but now also include two further points that we extracted from the




FIG. 16.—From KB99, isophote center coordinates X along the line joining P1 and P2 as a function of isophote major-axis radius a . A V -band *HST* WFPC2 image was measured twice, once masking out P2 (*green*) to measure the convergence of the P1 isophotes on the center of P1 (*green filled circle*) and once masking out P1 (*blue*) to measure the convergence of the P2+P3 isophotes on the center of P3 (*blue filled circle*). The brown and red symbols are measurements of individual isophotes in H - and K' -band images. The positions of the velocity center and of the center of mass of the BH and nucleus (if $M_{\bullet} = 3 \times 10^7 M_{\odot}$), each with error bars, are shown by the symbols labeled “ $V = 0$ ” and “COM,” respectively. The dashed line at $X = 0$ marks the center position of the bulge that was adopted by KB99. It was estimated by averaging all isophote center coordinates at $2''.9 < a < 25''.0$ ($1.3 < a^{1/4} < 2.24$). However, if $M_{\bullet} = 1.4 \times 10^8 M_{\odot}$, then the BH’s radius of influence is $r_{\text{cusp}} \simeq 7''.2$. Therefore, KB99 estimated the bulge center position partly from isophotes that are at $a < r_{\text{cusp}}$. Within this radius, the BH dominates the potential and isophotes do not need to be concentric to be in equilibrium (witness the eccentric disk). Since we now believe that the BH mass is large, we should derive the bulge center from correspondingly larger radii. The solid line is a least-squares fit to the bulge X_{center} values at $a > r_{\text{cusp}}$. It shows that the isophote centers at the largest radii in the figure are approximately at the X -coordinate of P3. Therefore, the BH is close to the luminosity-weighted center of the bulge.

QUIRC H -band image beyond this radius. In addition, we omit all center coordinates with errors larger than $0''.2$.

A least-squares fit to the points with $a > 7''.2$ gives the short black line in Figure 16. It shows that the bulge isophote centers drift with increasing radius toward the X -position of P3. Thus, the luminosity-weighted center of the bulge is close to P3.

This discussion suggests that the determination of the bulge center is less reliable than KB99 assumed. There are three reasons: (1) the BH sphere of influence is much larger than KB99 assumed, (2) the bulge isophote centers drift toward P3 with increasing radius beyond $a = 2''.9$, and (3) the isophote centers oscillate (or at least fluctuate) with radius because of dust or surface brightness fluctuations or perhaps a physical effect that we have not identified. If the BH is much more massive than P1+P2, then it is so close to the center of mass that M_{\bullet} cannot be determined accurately from the center-of-mass argument.

The important conclusion, however, is that the observations of M31 are consistent with dynamical equilibrium and with a large BH mass of $M_{\bullet} = 1.4 \times 10^8 M_{\odot}$.

8. ASTROPHYSICAL CONSTRAINTS ON A MASSIVE DARK OBJECT MADE OF DARK STARS

Central dark masses are detected dynamically in 38 galaxies (for reviews see Kormendy & Gebhardt 2001; Kormendy 2004).

¹⁷ What we should expect at $r < r_{\text{cusp}}$ is not clear. Because the potential is dominated by the BH, asymmetries like those of the P1-P2 eccentric disk are possible and isophotes do not need to be concentric to represent an equilibrium configuration.

They are commonly assumed to be supermassive BHs, although clusters of dark stars are consistent with the dynamics in most galaxies. Justifying this assumption, many authors cite the implausibility of producing so many stellar remnants (often 100 times the mass in visible stars) in the small volume defined by the PSF in which the dark mass must lie. Another argument is the consistency of the dark masses with energy requirements for BHs to power AGNs. More rigorous arguments against dark clusters are available for two galaxies, NGC 4258 and our own Galaxy (Maoz 1995, 1998; Genzel et al. 1998; Schödel et al. 2002, 2003; Ghez et al. 2005). Clusters of failed stars are not viable because brown dwarfs collide on short timescales and either evaporate or merge and become visible stars. Clusters of dead stars are not viable because their two-body relaxation times are so short that they evaporate. In NGC 4258, the timescales associated with these processes are at least as short as $10^{8.5}$ yr. In our own Galaxy, they are remarkably short indeed, $\sim 10^4$ yr. Even balls of neutrinos with cosmologically allowable neutrino masses are excluded in our Galaxy. The BH cases in NGC 4258 and in our Galaxy are now very strong and are taken as indications that dynamically detected central dark masses in other galaxies are BHs, too.

However, a great deal is at stake. It would be very important if astrophysical arguments ruled out BH alternatives in more than two galaxies. M32 has been the next best case (van der Marel et al. 1997, 1998), but Maoz (1998, Fig. 1) shows that a white dwarf cluster could survive for $\sim 10^{11}$ yr.

Applying our results on the dynamics of P3, M31 becomes the third galaxy in which dark star cluster alternatives to a BH can be excluded. For the most conservative estimate that $M_\bullet \simeq 3 \times 10^7 M_\odot$, the arguments are discussed in Kormendy et al. (2002) and Kormendy (2004). Here we update these arguments to the best kinematic fits and resulting BH masses implied by §§ 6 and 7. More detail is given in J. Kormendy et al. (2005, in preparation).

8.1. Limits on the Size of a Dark Cluster Alternative to a BH

Figure 17 derives our adopted limit on the size of any dark cluster alternative to a BH. It shows χ^2 contours for fits to the rotation and dispersion profiles of P3 using the thin-disk model of Figure 11 that gave the lowest BH mass. As in Maoz (1995, 1998), we assume that the dark object is a Plummer sphere, i.e., a reasonably realistic dynamical model with a very steep outer profile. We wish to use a relatively truncated mass distribution, one that is not excessively core-halo, because we need to fit the rapid rise in $V(r)$ and the corresponding drop in $\sigma(r)$ (Fig. 13) with a distributed object. Dark mass that is at several times the half-mass radius r_h hurts rather than helps us to do this.

We need to know how large r_h can be and still allow an adequate fit to the kinematics. As r_h is increased, the inner rotation curve drops, and it gets more difficult to fit the high central σ and especially the rapid central rise in $V(r)$. To compensate, an adequate fit requires that we increase M_\bullet . Therefore, r_h and M_\bullet are coupled (Fig. 17). How extended can the dark object be? We adopt the parameters at the upper right extreme of the 68% χ^2 contour: $r_h = 0''.031 = 0.113$ pc and $M_\bullet = 2.15 \times 10^8 M_\odot$. Note that choosing values corresponding to, e.g., the 90% χ^2 contour (or a larger one) does not significantly weaken the arguments against a dark cluster presented below because with increasing r_h the required dark cluster mass increases as well.

8.2. Arguments Against a Dark Cluster

The half-mass radius $r_h = 0.113$ pc is the same as the radius of the Ring Nebula (Cox 2000), a typical planetary nebula. We are




FIG. 17.—Contours of χ^2 for fits to the P3 kinematics of Plummer spheres with half-mass radii r_h and total masses M_\bullet . The P3 model is a flat disk, as in Fig. 11. Our adopted constraint on the fluffiness of the dark mass is the upper right extreme of the 68% χ^2 contour, i.e., $r_h = 0''.031 = 0.113$ pc and $M_\bullet = 2.15 \times 10^8 M_\odot$.

considering a situation in which this volume contains $10^8 M_\odot$ of brown dwarf stars or stellar remnants. The mean density inside r_h is $\rho_h = 1.8 \times 10^{10} M_\odot \text{pc}^{-3}$, and the density at r_h is $\rho(r_h) = 6.5 \times 10^9 M_\odot \text{pc}^{-3}$. This is ~ 10 times larger than the largest stellar mass density observed in any galaxy, $7 \times 10^8 M_\odot \text{pc}^{-3}$ at $r = 0''.1 = 0.004$ pc in the stellar cusp around Sgr A* in our Galaxy (Genzel et al. 2003). However, only about $300 M_\odot$ of stars are inside the above radius (Genzel et al. 2003). Not surprisingly, a dark cluster as extreme as the one that we require to explain the kinematics of P3 gets into trouble.

8.2.1. Brown Dwarfs Collide and Destroy Themselves

It is easiest to eliminate brown dwarfs. They collide with each other so violently that they get converted back into gas. Figure 18 shows the timescale on which every typical brown dwarf collides with another brown dwarf. As in Maoz (1995, 1998), the zero-temperature brown dwarf radius is taken from Zepolsky & Salpeter (1969) and Stevenson (1991), and the calculation is an average interior to r_h . Typical collision velocities at r_h are $\sim 2500 \text{ km s}^{-1}$; this is fast enough compared to the surface escape velocity ($\sim 600 \text{ km s}^{-1}$ for a $0.08 M_\odot$ star and smaller for lower mass brown dwarfs) that the brown dwarfs would get destroyed, i.e., converted back into gas. Brown dwarfs are strongly excluded.

8.2.2. Intermediate-Mass White Dwarfs Collide and Make Type Ia Supernovae

Relatively short collision times provide an argument against intermediate-mass white dwarfs. For $0.8 M_\odot \lesssim m_* \lesssim 1.2 M_\odot$, collision times at the quarter-mass radius $r_{1/4}$ are $(4\text{--}7) \times 10^9$ yr. Given the implied numbers of white dwarfs interior to this radius and the fact that the collision time would be shorter at smaller radii, collisions should happen more often than every 50–150 yr. Each collision would bring the remnant well above the Chandrasekhar limit. Presumably Type Ia supernovae would result. Near maximum brightness, they would be visible to the naked eye. The fact that no such supernovae have been observed in M31 might barely




FIG. 18.—Timescales on which dark cluster alternatives to a BH get into trouble in M31. The dark star mass is m_* . For a cluster made of brown dwarfs (“BD”), the left line shows the timescales on which every typical star suffers a physical collision with another star. Points WD, NS, and BH show the times in which dark clusters made of white dwarfs, neutron stars, or stellar-mass BHs would evaporate. Points P with “error bars” are the timescales on which every typical progenitor star collides with another progenitor at the radius in the Plummer model dark cluster that contains one-quarter of the total mass. The letter P is for the time when the dark cluster is three-quarters assembled; the “error bars” end at the collision times when the cluster is half assembled (top) and fully assembled (bottom).

be consistent with the above rates, but if intermediate-mass white dwarfs in similar dark clusters are the explanation for other galaxies’ central dark objects, the resulting supernovae would easily have been seen. Intermediate-mass white dwarfs are implausible.

In addition, the supernova ejecta would be lost to the cluster. The above collision times imply that most of the mass inside $r_{1/4}$ and a significant fraction of the mass inside r_h would be lost in a few billion years. For the dark cluster to have its present mass, it would have had to be more massive in the past. All problems involving collision rates would get more severe.

White dwarfs with masses less than half of the Chandrasekhar limit will turn out to be excluded because their progenitors would be destroyed and converted into gas or, if they succeed to merge, become progenitors of intermediate-mass white dwarfs or still heavier remnants (§ 8.2.4).

White dwarfs with masses near the Chandrasekhar limit are small. Their collision times are long. For these objects, we need stronger arguments. These arguments will militate against intermediate-mass white dwarfs, also.

8.2.3. Dark Cluster Formation Scenario: Let’s Imagine Six Impossible Things before Breakfast

Heavy remnants are too small to collide. Instead, relaxation gives positive energies to a steady trickle of stars that are lost to the system. In $\lesssim 300$ half-mass relaxation times, the cluster evaporates. Figure 18 shows the evaporation times for dark clusters made of $0.6 M_\odot$ white dwarfs, $0.8 M_\odot$ white dwarfs, $1.2 M_\odot$ white dwarfs, $1.5 M_\odot$ neutron stars, and $3 M_\odot$ BHs (left to right; symbols WD, WD, WD, NS, and BH). Unlike the case in NGC 4258 and our Galaxy (Maoz 1995, 1998), these evaporation times are not

implausibly short except for $m_* \gtrsim 10 M_\odot$ BHs. So, for most remnants, we need stronger arguments.

Fortunately, we can add new arguments. They depend only on canonical, well-understood stellar evolution and simple stellar dynamics. A dark cluster made of stellar remnants is viable only if its progenitor stars can safely live their lives and deliver their remnants at suitable radii. The properties of the dark cluster constrain how it can form. We describe the most benign formation scenario in this subsection. It requires fine-tuning of the star formation in ways that we do not know are possible. However, we do not base our arguments against the resulting dark clusters on these problems, because we do not understand star formation well enough. But main-sequence stars are well understood, and we know progenitor star masses well enough for the present purposes. It turns out that progenitor stars get into trouble because they must be so close together that they collide. The consequences are untenable, as discussed in the following subsections.

Finding a plausible formation scenario is comparable to imagining six impossible things before breakfast. The argument is summarized as follows. The progenitor cluster must be as small as the dark cluster because dynamical friction is too slow to deliver remnants from much larger radii. From Lauer et al. (1998), the density of P2 at $r \sim 0.1\text{--}0.2$ pc is $\sim 10^6 M_\odot \text{pc}^{-3}$. Then the characteristic time for dynamical friction (Binney & Tremaine 1987, eq. [7-18]) to change velocities $v \sim 10^3 \text{ km s}^{-1}$ is $v/(dv/dt) \sim 10^{12}$ yr for $10 M_\odot$ stars. This drives us to imagine the following impossible things:

1. Let us form progenitor stars with a density distribution proportional to that of the dark cluster, i.e., a Plummer sphere with half-light radius $r_h = 0.113$ pc.
2. We get into less trouble with collisions if fewer progenitors are resident at one time. Therefore, the safest strategy is to form stars at a constant rate during the formation time of, say, 10^{10} yr. This is not the obvious strategy in a hierarchically clustering universe; it is more natural to postulate episodic formation by more vigorous events that are connected with major mergers. But shortening the formation time increases the number of progenitors that must be resident at the same time, and this greatly increases difficulties with stellar collisions.
3. We assume that all progenitor stars have the same mass. In particular, we cannot allow a Salpeter (1955) mass function because we cannot tolerate any significant numbers of dwarf stars with lifetimes long enough so that the stars or their white dwarf remnants remain visible today.
4. We assume that sufficient gas for star formation is always present. Some gas could come from mass lost by evolving stars, but some gas must be added continuously to make the cluster grow. We assume that stars can form despite any energy feedback from massive or evolved stars.
5. We do not worry about the fact that the young cluster is easy to unbind gravitationally by the mass loss from evolved stars. This is a difficult problem. Progenitors outmass their remnants by factors of at least a few (for low-mass stars) or ~ 10 (for high-mass stars). During the first stellar generations, the progenitors outmass the remnants. Since they lose most of their mass during the course of stellar evolution, it is easy to reduce the total mass of the cluster substantially when stars die. Impulsive loss of more than half of the mass (say, if the star formation happened in a coeval starburst) unbinds the cluster. Slower mass loss fights the formation process by expanding the cluster. We ignore all of these difficulties and assume that the cluster can safely evolve beyond the fragile initial stage when the mass of progenitors present at one time is significant.

6. We assume that the only evolution in r_h is that resulting from a gradual increase of the cluster mass. Then $r_h \propto M_\bullet^{-1}$.

Using the above assumptions, we calculate the evolution of the cluster for various combinations of progenitors and their remnants. Progenitor masses are from Iben et al. (1996) for white dwarfs and from Brown & Bethe (1994) for BHs. The progenitor clusters get into the following trouble.

8.2.4. *If M31 Is Typical, Then Progenitor Clusters Are Too Bright*

The above progenitor clusters have absolute magnitudes ranging from $M_V \simeq -16.3$ to -17.5 for the duration of their formation. These absolute magnitudes are almost independent of progenitor star mass; higher mass progenitors are much more luminous, but they live much less long, so far fewer are present at one time. Nuclei as bright as the above could not be hidden in nearby (or even moderately distant) galaxies. They are rare (e.g., Lauer et al. 1995, 2005). It is unreasonable to assume that dark cluster formation lasted for $\sim 10^{10}$ yr in every bulge and then stopped recently in all galaxies.

If the formation of the dark cluster took $\ll 10^{10}$ yr, then the progenitor clusters are brighter but it is easier to hide them at large redshifts. But then all problems that involve stellar collisions get much worse (see below).

This problem applies to all types of stellar remnants.

8.2.5. *Dynamical Friction Deposits Remnants at Small Radii*

As noted above, progenitor stars are much more massive than the remnants of previous generations that already make up the dark cluster. The dynamical friction of the progenitors against the remnants makes the progenitors sink quickly to small enough radii so that the progenitor cluster becomes self-gravitating. Two problems result. Progenitor collision times get shorter; these are discussed further below. Second, remnants are deposited at small radii, inconsistent with the density distribution that we are trying to construct. As heavy stars sink, remnants are lifted to higher radii; the effect is not large for one generation of progenitors, but it adds up by the time the cluster is finished. The result is a dark cluster that is much more core-halo in structure than a Plummer model. That is, it is inherently impossible to make a dark cluster that is as centrally concentrated as a Plummer model via progenitor stars that greatly outmass their remnants. This is one of the stiffest problems of our formation scenario.

If the dark cluster is less compact than a Plummer sphere, then its half-mass radius must be smaller than 0.113 pc in order to fit the kinematics. All problems with stellar collisions get much worse.

This problem also applies to all types of remnants.

8.2.6. *White Dwarfs that Cannot Merge to Form Type Ia Supernovae Cannot Be Relevant*

Interior to r_h , essentially all progenitors of $0.6 M_\odot$ white dwarfs collide and either get destroyed (their surface escape velocities are $\sim 600 \text{ km s}^{-1}$) or, in the earlier phases of the dark cluster formation, merge. If they merge, they get converted to progenitors of white dwarfs that have masses $\gtrsim 0.8 M_\odot$. Progenitors of white dwarfs with masses $\lesssim 0.55 M_\odot$ live too long to have died and, provided that they were not destroyed, would still be visible. Therefore, white dwarfs that are low enough in mass so that a collision of two of them results in a remnant that is less massive than the Chandrasekhar limit are not relevant.

8.2.7. *Progenitors Collide and Evaporate or Merge into High-Mass Stars*

Figure 18 shows progenitor star collision times for the second half of the dark cluster formation process. At $r \lesssim r_{1/4}$,

progenitors of low-mass remnants get destroyed and converted into gas or, in earlier phases, merge to become progenitors of high-mass remnants. We note that the stellar evolution clock is reset to essentially zero in every nondestructive collision because the merging stars get thoroughly mixed. Dynamical friction is neglected in constructing Figure 18; if it is included, then most of the progenitors participate in the collisions. Also neglected is the fact that successive mergers increase the mass range and hence decrease both the dynamical friction sinking time and the relaxation time of the cluster.

Three consequences spell trouble for the formation scenario:

1. If progenitors are not destroyed, they merge up to form stars of high enough masses so that they die as Type II supernovae. Their luminosity is not a problem for the hypothetical, present M31 dark cluster because its formation process is finished or at least in hiatus. But again, if M31 is typical, then the formation of many such objects at intermediate and high redshifts should produce one Type II supernova per galaxy per ~ 100 yr at the center of the galaxy. They would have been seen.

2. The supernova ejecta again would not be bound to the dark cluster unless a large amount of gas is also present.

3. Relatively few, high-mass remnants would be formed. Dynamical friction would guarantee that they got deposited at small radii. The mass range that resulted from heterogeneous stellar merger histories would create a large mass range even if none was present initially. The result would be that relaxation times would be much shorter, plausibly an order of magnitude shorter, than the single-mass relaxation times that gave rise to the cluster evaporation times in Figure 18. For all of these reasons, evaporation times are likely to be much shorter than the several billion years indicated for $3 M_\odot$ BHs in Figure 18. This is implausibly short.

8.2.8. *Summary on BH Alternatives*

Therefore, astrophysically reasonable alternatives to a supermassive BH are likely to fail. The arguments against brown dwarfs seem bomb proof. The arguments against stellar remnants are more complicated, but they are based on secure aspects of stellar and star-cluster evolution. Also, there are many arguments, even a few of which are sufficient. So we are not very vulnerable to uncertainties involving any one argument (“Are we sure that we have not missed those supernovae or confused them with AGN activity?”). The problem (§ 8.2.5) that remnants are deposited at excessively small radii is particularly important. In this paper we have derived the largest published estimate of M_\bullet using data at the smallest radii. This leaves little room for distributed dark matter, i.e., for a dark cluster with core-halo structure. In addition, § 8.2.3 on the formation scenario, while not formally part of our argument against dark clusters, presents formidable challenges. Our arguments are discussed in more detail in J. Kormendy et al. (2005, in preparation). However, our conclusion that dark cluster alternatives to a BH are excluded seems robust.

9. CONCLUSIONS

M31 is now the third galaxy in which astrophysical arguments strongly favor the conclusion that a dynamically detected central dark object is a BH. M31 is the only galaxy for which such arguments are based on *HST* observations. Similar conclusions for NGC 4258 and our Galaxy result from ground-based observations. The present result is therefore an important contribution of *HST* to the BH paradigm of AGNs. It increases our confidence that all dynamically detected central dark objects in galaxies are BHs.

We are most grateful to Don Winget for very helpful discussions about white dwarf properties and to Don Winget and Anjum Mukadam for providing digital observed spectra of white dwarf stars. We thank Detlev Koester for making his models of white dwarf atmospheres available and for insightful remarks about their systematics. We also thank the Nuker team (D. Richstone, PI) and especially Scott Tremaine for many helpful and enlightening discussions. R. B. is most grateful to the Department of Astronomy of the University of Texas at Austin for its warm hospitality and the support provided by a Beatrice M. Tinsley Centennial Visiting Professorship. We thank the anonymous referee for a careful reading of the

paper and for very helpful suggestions that led to significant improvements in §§ 1 and 8 and that prompted us to add § 6.2.

This paper is based on observations made with the NASA/ESA *Hubble Space Telescope*, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. These observations are associated with program 8018 (visits 1 and 2) and were a portion of time allocated to the Space Telescope Imaging Spectrograph Instrument Definition Team key project for supermassive BHs. *Facilities*: HST(STIS)

APPENDIX

Peiris & Tremaine (2003) show that their eccentric disk model fitted to ground-based kinematic data also agrees remarkably well with our STIS kinematic measurements of P1 and P2. The comparison includes not only V and σ but also the parameters h_3 and h_4 , which measure the lowest order departures from Gaussian line profiles. The data that they use are presented here in Tables 4 and 5 and Figures 19 and 20. We do not repeat their discussion. Instead, we focus on the generic properties of the LOSVDs. In particular, we confirm an unusual property of the LOSVDs that directly implies aligned, eccentric orbits. This effect was seen and interpreted in KB99, but it is much larger at *HST* spatial resolution. Since the effect was inherent in but not explicitly predicted by Tremaine (1995), it is compelling evidence in favor of his model.

Figure 19 compares the FQ and FCQ reductions of the Ca infrared triplet spectra. The velocities are almost identical. The dispersions agree where the higher order Gauss-Hermite coefficients are small and differ in the expected way when they are not. Where FCQ measures an extended wing of the LOSVD in the prograde direction ($h_3 < 0$ at $r \simeq -0''.1$), it finds a smaller dispersion than FQ because FQ fits a Gaussian to the whole LOSVD, including wings. Similarly, where $h_4 > 0$ at $r \simeq +0''.1$, FCQ fits a broader Gaussian than FQ and then clobbers the intermediate-velocity shoulders of the Gaussian with h_4 to make the fitted LOSVD triangular.

Kinematic asymmetries inherent in an eccentric disk provided the basic test of that model in Tremaine (1995), KB99, and Peiris & Tremaine (2003). We now know that the BH is in the blue nucleus P3 at $r = 0$ in Figure 19. Stars in the eccentric disk linger at apocenter to form P1; as a result, both the rotation amplitude $|V|$ and the velocity dispersion σ are relatively small at $r \simeq +0''.5$. Other stars in the eccentric disk are passing pericenter slightly on the anti-P1 side of the BH; as a result, the rotation amplitude is large at $r \simeq -0''.2$ in P2. The apparent velocity dispersion is highest at approximately the same radii because the slit and PSF average over stars moving in a variety of directions as they swing around the BH. Figures 11 and 12 in Peiris & Tremaine (2003) show that their nonaligned model accurately accounts for the asymmetric rotation and dispersion profiles.

TABLE 4
KINEMATICS OF M31 DERIVED FROM THE RED CAT SPECTRA WITH THE FOURIER QUOTIENT METHOD

Radius (arcsec)	V (km s ⁻¹)	ΔV (km s ⁻¹)	σ (km s ⁻¹)	$\Delta\sigma$ (km s ⁻¹)
-1.075	-177	14	130	15
-0.726	-189	13	151	14
-0.474	-244	13	179	14
-0.304	-299	23	207	24
-0.230	-319	32	274	35
-0.180	-332	34	298	38
-0.130	-276	36	345	39
-0.080	-232	45	373	48
-0.005	-118	23	247	25
0.070	-52	28	221	31
0.120	0	22	195	23
0.170	32	23	220	26
0.220	84	21	182	23
0.270	108	14	174	15
0.320	140	15	186	17
0.396	175	10	146	11
0.495	207	8	131	8
0.595	192	7	122	8
0.694	200	7	117	8
0.794	200	7	104	8
0.894	185	9	103	11
0.994	178	11	111	13
1.094	156	10	91	12
1.262	175	11	99	13
1.513	147	19	108	23

TABLE 5
KINEMATICS OF M31 DERIVED FROM THE RED CAT SPECTRA WITH THE FOURIER CORRELATION QUOTIENT METHOD

Radius (arcsec)	V (km s ⁻¹)	ΔV (km s ⁻¹)	σ (km s ⁻¹)	$\Delta\sigma$ (km s ⁻¹)	h_3	Δh_3	h_4	Δh_4
-1.390	-130.8	11.7	135.4	10.6	0.040	0.079	-0.092	0.079
-1.023	-189.6	11.0	140.0	11.6	0.056	0.072	-0.043	0.072
-0.776	-192.2	8.3	144.2	10.2	0.110	0.053	0.013	0.053
-0.578	-217.0	11.2	188.7	11.8	0.037	0.054	-0.046	0.054
-0.401	-267.2	7.2	168.3	7.1	0.168	0.039	-0.070	0.039
-0.254	-316.4	12.7	240.2	12.2	0.082	0.048	-0.074	0.048
-0.180	-313.6	14.4	279.6	15.5	-0.089	0.047	-0.035	0.047
-0.130	-247.9	16.3	296.5	22.1	-0.188	0.050	0.060	0.050
-0.080	-181.4	16.1	300.7	23.1	-0.264	0.049	0.084	0.049
-0.030	-137.6	14.5	273.7	18.2	-0.147	0.048	0.026	0.048
0.020	-99.0	13.0	263.9	17.3	-0.156	0.045	0.051	0.045
0.070	-59.6	15.6	286.1	27.5	-0.061	0.050	0.194	0.050
0.120	-4.8	12.6	226.0	21.5	-0.072	0.051	0.176	0.051
0.170	36.0	13.0	231.1	18.4	-0.001	0.051	0.079	0.051
0.246	96.2	8.4	184.9	11.5	0.009	0.041	0.062	0.041
0.346	152.2	7.0	170.3	8.2	-0.042	0.038	-0.007	0.038
0.446	198.2	4.8	143.3	5.7	-0.075	0.031	-0.002	0.031
0.592	203.6	3.5	126.1	3.9	-0.072	0.025	-0.021	0.025
0.788	199.2	3.5	109.2	3.8	-0.016	0.029	-0.029	0.029
0.989	185.5	5.7	117.0	6.1	-0.185	0.045	-0.037	0.045
1.213	173.6	5.9	103.1	8.2	0.014	0.052	0.075	0.052
1.569	140.3	7.0	108.7	12.5	-0.379	0.059	0.221	0.059

An important additional property of the LOSVDs (and the unusual one mentioned above) is the observation that h_3 has the same sign as V over a radius range of $\Delta r \simeq 0''.4$ centered $\sim 0''.1$ on the anti-P1 side of P3. This effect was seen at ground-based spatial resolution in KB99. It is much larger here. It is opposite to the normal behavior of rotating stellar systems, in which velocity projection along the line of sight makes h_3 antisymmetric with V (van der Marel et al. 1994; Bender et al. 1994). Also, the maximum amplitude, $h_3 \simeq -0.3$, is unusually large compared to values in other galaxies. All this is easily seen in the LOSVDs (Fig. 20; radii $-0''.15$ and $-0''.10$) as the broad wings on the $-V$ side of the line centers. These wings tell us that, on the anti-P1 side of the P3, where the average galactic rotation is toward us (V is negative in Fig. 20), many stars are rotating more rapidly and few stars are rotating more slowly than the mean rotation velocity averaged within the PSF.




FIG. 19.—Red filled circles: Rotation velocity V , velocity dispersion σ , and Gauss-Hermite parameters h_3 and h_4 as a function of radius as derived with FCQ from the red spectrum of the nucleus of M31. Blue filled circles: FQ results from Fig. 2.




FIG. 20.—LOSVDs in the same radial bins that were used to derive the FCQ kinematic results shown in Fig. 19. The black lines are nonparametric LOSVDs; the red lines are the FCQ fits. The radius of the bin is given at upper right in each panel. At upper left, we tabulate the velocity dispersion and the Gauss-Hermite parameters h_3 and h_4 .

Our interpretation is the same as in KB99. The velocity dispersion is expected to look big near pericenter in the eccentric disk because the slit and PSF integrate over stars that are at different positions along orbits that are rapidly turning around the BH. A prograde LOSVD wing follows naturally if there are many stars still closer to the BH and if they also are in eccentric orbits with apocenters that point toward P1. In the almost Keplerian potential of the BH, these stars have larger pericenter velocities than the mean V farther out; in fact, their velocities should be larger than the local circular velocity. Consistent with this interpretation, the LOSVD asymmetry is most obvious at $r = -0''.05$ to $-0''.15$, i.e., at slightly more than one PSF radius on the anti-P1 side of P3. The highest velocities reach $\sim 1000 \text{ km s}^{-1}$, indeed somewhat larger than what we measure at about the same location for the PSF-blurred velocities of the blue stars in P3. Of course, this explanation only works if the BH is embedded in P3.

The fact that we can understand naturally an observation not predicted by Tremaine (1995) increases our confidence in his model. With improved disk parameters, Peiris & Tremaine (2003, see Figs. 13 and 14) accurately predict the h_3 and h_4 profiles near the BH. At this point, there seems little doubt that the interpretation of the double nucleus as an eccentric disk is correct and that its main parameters have been determined. The important next step is self-consistent dynamical models to investigate whether the present configuration can be long lived (e.g., Statler et al. 1999; Statler 1999; Bacon et al. 2001; Salow & Statler 2004). A larger BH mass, as estimated here, will likely help to construct more long-lived models. Beyond that, the origin of the eccentric disk remains essentially unknown.

REFERENCES

- Aparicio, A., & Gallart, C. 2004, *AJ*, 128, 1465
 Bacon, R., Emsellem, E., Monnet, G., & Nieto, J.-L. 1994, *A&A*, 281, 691
 Bacon, R., et al. 2001, *A&A*, 371, 409
 Bender, R. 1990, *A&A*, 229, 441
 Bender, R., Saglia, R. P., & Gerhard, O. E. 1994, *MNRAS*, 269, 785
 Bertola, F., Bressan, A., Burstein, D., Buson, L. M., Chiosi, C., & di Serego Alighieri, S. 1995, *ApJ*, 438, 680
 Binney, J., & Tremaine, S. 1987, *Galactic Dynamics* (Princeton: Princeton Univ. Press)
 Bower, G. A., et al. 2001, *ApJ*, 550, 75
 Brown, G. E., & Bethe, H. A. 1994, *ApJ*, 423, 659
 Brown, T. M., Ferguson, H. C., Stanford, S. A., & Deharveng, J.-M. 1998, *ApJ*, 504, 113
 Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 1000
 Corbin, M. R., O'Neil, E., & Rieke, M. J. 2001, *AJ*, 121, 2549
 Cox, A. N. 2000, *Allen's Astrophysical Quantities* (New York: Springer)
 Crane, P., et al. 1993a, *AJ*, 106, 1371
 Crane, P. C., Cowan, J. J., Dickel, J. R., & Roberts, D. A. 1993b, *ApJ*, 417, L61
 Crane, P. C., Dickel, J. R., & Cowan, J. J. 1992, *ApJ*, 390, L9
 Davidge, T. J., Rigaut, F., Doyon, R., & Crampton, D. 1997, *AJ*, 113, 2094
 Dressler, A. 1984, *ApJ*, 286, 97
 Dressler, A., & Richstone, D. O. 1988, *ApJ*, 324, 701
 Emsellem, E., & Combes, F. 1997, *A&A*, 323, 674
 Ferrarese, L., & Merritt, D. 2000, *ApJ*, 539, L9
 Finley, D. S., Koester, D., & Basri, G. 1997, *ApJ*, 488, 375
 Gebhardt, K., et al. 2000a, *AJ*, 119, 1157
 ———. 2000b, *ApJ*, 539, L13
 ———. 2003, *ApJ*, 583, 92

- Genzel, R., et al. 1998, *ApJ*, 498, 579
———. 2003, *ApJ*, 594, 812
- Ghez, A. M., Salim, S., Hornstein, S. D., Tanner, A., Lu, J. R., Morris, M., Becklin, E. E., & Duchêne, G. 2005, *ApJ*, 620, 744
- Ghez, A. M., et al. 2003, *ApJ*, 586, L127
- Goudfrooij, P., Baum, S. A., & Walsh, J. R. 1997, in *The 1997 HST Calibration Workshop*, ed. S. Casertano et al. (Baltimore: STScI), 100
- Iben, I., Tutukov, A. V., & Yungelson, L. R. 1996, *ApJ*, 456, 750
- Jacobs, V., & Sellwood, J. A. 2001, *ApJ*, 555, L25
- King, I. R., Stanford, S. A., & Crane, P. 1995, *AJ*, 109, 164
- King, I. R., et al. 1992, *ApJ*, 397, L35
- Kleinman, S. J., et al. 2004, *ApJ*, 607, 426
- Koester, D., et al. 2001, *A&A*, 378, 556
- Kormendy, J. 1987, in *IAU Symp. 127, Structure and Dynamics of Elliptical Galaxies*, ed. T. de Zeeuw (Dordrecht: Reidel), 17
———. 1988, *ApJ*, 325, 128
———. 2004, in *Coevolution of Black Holes and Galaxies*, ed. L. C. Ho (Cambridge: Cambridge Univ. Press), 1
- Kormendy, J., & Bender, R. 1999, *ApJ*, 522, 772 (KB99)
- Kormendy, J., Bender, R., & Bower, G. 2002, in *ASP Conf. Ser. 273, The Dynamics, Structure and History of Galaxies*, ed. G. S. Da Costa & H. Jerjen (San Francisco: ASP), 29
- Kormendy, J., & Gebhardt, K. 2001, in *AIP Conf. Proc. 586, 20th Texas Symposium on Relativistic Astrophysics*, ed. J. C. Wheeler & H. Martel (Melville: AIP), 363
- Kormendy, J., & Kennicutt, R. C., Jr. 2004, *ARA&A*, 42, 603
- Lauer, T. R., et al. 1993, *AJ*, 106, 1436
———. 1995, *AJ*, 110, 2622
———. 1998, *AJ*, 116, 2263
———. 2005, *AJ*, 129, 2138
- Le Borgne, J.-F., et al. 2003, *A&A*, 402, 433
- Light, E. S., Danielson, R. E., & Schwarzschild, M. 1974, *ApJ*, 194, 257
- Magorrian, J., et al. 1998, *AJ*, 115, 2285
- Maoz, E. 1995, *ApJ*, 447, L91
———. 1998, *ApJ*, 494, L181
- Morris, M. 1993, *ApJ*, 408, 496
- Mould, J., Graham, J., Matthews, K., Soifer, B. T., & Phinney, E. S. 1989, *ApJ*, 339, L21
- Nieto, J.-L., Macchetto, F. D., Perryman, M. A. C., di Serego Alighieri, S., & Lelièvre, G. 1986, *A&A*, 165, 189
- Peiris, H. V., & Tremaine, S. 2003, *ApJ*, 599, 237
- Pickles, A. J. 1998, *PASP*, 110, 863
- Rich, R. M., Mighell, K. J., Neill, J. D., & Freedman, W. L. 1996, in *New Extragalactic Perspectives in the New South Africa*, ed. D. L. Block & J. M. Greenberg (Dordrecht: Kluwer), 325
- Richstone, D., Bower, G., & Dressler, A. 1990, *ApJ*, 353, 118
- Salow, R. M., & Statler, T. S. 2001, *ApJ*, 551, L49
———. 2004, *ApJ*, 611, 245
- Salpeter, E. E. 1955, *ApJ*, 121, 161
- Sambhus, N., & Sridhar, S. 2002, *A&A*, 388, 766
- Sanders, R. H. 1998, *MNRAS*, 294, 35
- Schmidt, M. 1959, *ApJ*, 129, 243
- Schödel, R., Ott, T., Genzel, R., Eckart, A., Mouawad, N., & Alexander, T. 2003, *ApJ*, 596, 1015
- Schödel, R., et al. 2002, *Nature*, 419, 694
- Schwarzschild, M. 1979, *ApJ*, 232, 236
- Sérsic, J. L. 1968, *Atlas de Galaxias Australes* (Cordoba: Observatorio Astronomico)
- Statler, T. S. 1999, *ApJ*, 524, L87
- Statler, T. S., King, I. R., Crane, P., & Jedrzejewski, R. I. 1999, *AJ*, 117, 894
- Stevenson, D. J. 1991, *ARA&A*, 29, 163
- Thomas, J., Saglia, R. P., Bender, R., Thomas, D., Gebhardt, K., Magorrian, J., & Richstone, D. 2004, *MNRAS*, 353, 391
- Tonry, J. L. 1984, *ApJ*, 283, L27
———. 1987, *ApJ*, 322, 632
- Tonry, J. L., et al. 2001, *ApJ*, 546, 681
- Toomre, A. 1964, *ApJ*, 139, 1217
- Tremaine, S. 1995, *AJ*, 110, 628
———. 2001, *AJ*, 121, 1776
- Tremaine, S., et al. 2002, *ApJ*, 574, 740
- van der Marel, R. P., Cretton, N., de Zeeuw, T., & Rix, H.-W. 1998, *ApJ*, 493, 613
- van der Marel, R. P., Rix, H.-W., Carter, D., Franx, M., White, S. D. M., & de Zeeuw, T. 1994, *MNRAS*, 268, 521
- van der Marel, R. P., et al. 1997, *Nature*, 385, 610
- Verolme, E. K., et al. 2002, *MNRAS*, 335, 517
- Yu, Q. 2003, *MNRAS*, 339, 189
- Zapolsky, H. S., & Salpeter, E. E. 1969, *ApJ*, 158, 809